



Italian folk plant-based remedies to heal headache (XIX-XX century)



Rosalucia Mazzei^a, Elvira V. De Marco^b, Olivier Gallo^b, Giuseppe Tagarelli^{a,*}

^a Institute for Agricultural and Forest Systems in the Mediterranean, National Research Council, Via Cavour 4-6, 87036 Rende (CS), Italy

^b Institute of Neurological Sciences, National Research Council, C.da Burga, 87050 Mangone (CS), Italy

ARTICLE INFO

Keywords:

Headache
Folk medicine
Plant remedies
Ethnomedicine
Plant-based drugs

ABSTRACT

Background: Headache has been recognized since antiquity. From the late nineteenth to the early to mid-twentieth century, Italian folk remedies to treat headache were documented in a vast *corpus* of literature sources. **Aim:** The purpose of this paper is to bring to light the plant-based treatments utilized by Italian folk medicine to heal headache in an attempt to discuss these remedies from a modern pharmacological point of view. Moreover, we compare the medical applications described by Hippocrates, Pliny the Elder, Dioscorides, Galen and Serenus Sammonicus with those utilized by Italian folk medicine to check if they result from a sort of continuity of use by over two thousand years.

Results: A detailed search of the scientific data banks such as Medline and Scopus was undertaken to uncover recent results concerning the anti-inflammatory, anti-nociceptive and analgesic activities of the plants. Fifty-eight (78.4%) plant-based remedies have shown *in vivo*, *in vitro* or in human trials a large spectrum of anti-inflammatory, anti-nociceptive and analgesic activities.

Moreover, thirty-one of remedies (41.9%) were already included in the pharmacopoeia between the 5th century BC and the 2nd century AD.

Conclusion: Italian folk medicine could be a promising source of knowledge and could provide evidences for active principles that have not as of yet been fully used for their potential.

1. Introduction

The third edition of the International Headache Society classifies headache into “primary” and “secondary” disorders. The primary headaches, representing 90% of all cephalalgias, include migraine, tension-type headache, cluster headache, trigeminal autonomic cephalalgias, and other primary headaches (such as cold-stimulus headache, external-pressure headache, etc). The secondary ones are caused by an underlying disease, such as head injuries, infections, vascular disorders or tumors (HIS, 2013). The World Health Organization claims that, nowadays, headache is among the most prevalent disorders in the world (WHO, 2011), but it is reasonable to state headache has existed as long as humankind (Magiorkinis et al., 2009).

In fact, in the opinion of several researchers, early medical treatments to relieve headache sufferers were cranial trepanations performed by Neolithic populations (around 9000 BCE). The practice of removing pieces of bone from skull of a living individual was connected with the belief to release the evil spirits inhabited in the head of the patients (Campillo, 1984). The idea that the cause of headache, as other

ailments, could be ascribable to an attack of divine or supernatural forces, was common during antiquity. The first written descriptions of headache symptoms were found in Mesopotamian tablets (4000 BCE) in which the attack was attributed to an evil called *Tíu*. The pain was cured with an exorcism using an ointment of human bone reduced to ashes and mixed with cedar oil or applying other unpleasant substances (Green et al., 2005).

Later, ancient Egyptian medical papyri, such as the Ebers papyrus (1550 BCE) contained remedies and prescriptions corresponding to therapeutic remedies which ranged from supernatural to natural, from magical to empirical (Karenberg and Leitz, 2001).

The beginning of a rational approach arises from Greek medicine: Hippocrates of Kos (460–377 BCE) argued that disease is a general phenomenon of organism without supernatural interventions. He suggested headache occurred when black bile, yellow bile, blood and phlegm were out of harmony in the body, so he treated head pain with bloodletting or applying herbs to the head to drain excess liquids. Hippocrates was, also, the first to describe a migraine with aura (Green et al., 2005).

Abbreviation: NSAIDs, Nonsteroidal anti-inflammatory drugs; NF-κB, Nuclear Factor-κB; COX-2, Cyclooxygenase-2; PGE-2, Prostaglandin E-2; TNF-α, Tumor Necrosis Factor-α; NO, Nitric Oxide; IL-1, Interleukin-1; IL-8, Interleukin-8; LPS, lipopolysaccharide; MAO-A, Monoamino Oxidase-A; 5-HT, 5-hydroxytryptamine; NE, Norepinephrine

* Corresponding author.

E-mail address: giuseppe.tagarelli@cnr.it (G. Tagarelli).

In the Roman period, Pliny the Elder (23–79 CE) who was not a physician but a writer, in his *Naturalis Historia*, cites about one hundred remedies taken from animal (man included) and vegetable kingdoms. He subdivides headaches into five categories, based on the symptoms: 1) pain of entire head; 2) pain localized to the temples; 3) pain to the nape; 4) pain accompanied by sense of warmth and, lastly, 5) chronic headache. The contemporaries Aulus Cornelius Celsus (c.25 BCE–c.50 CE) and Aretus the Cappadocian (1st century AD) recommend, as therapeutic remedies, bloodletting, diet, application of warm or cold water and cauterization. Later, Galen (129–199 CE) was the first to introduce the term *hemicrania* (Zanchin, 2010).

The purpose of this paper is to bring to light the plant-based treatments utilized by Italian folk medicine for headache, between the late nineteenth century and the early to mid-twentieth century, to “rediscover” remedies that nowadays are mostly no longer used. For this reason, a systematic review of literature sources was carried out to investigate practices, performed by Italian folk medicine to heal headache during the above mentioned period, in the light of the current scientific knowledge.

The starting point of this period stands for the time in which the study of folk traditions as folk treatments to heal several diseases, arose and spread out with an appropriate methodology (Cirese, 1996). On the other hand, the end of the 1950's represents the period in which the countryside was massively abandoned so that, for the first time in Italy, more people were employed in industry than in agriculture (ISTAT, 2011). This event played an important role in the so called “economic boom” and in the social and cultural change of Italy which was transformed from a mainly rural to a modern country (Sapelli, 1991).

Moreover, in this paper we compare medical applications described by Hippocrates, Pliny the Elder, Dioscorides, Galen and Serenus Sammonicus to verify possible similarities with Italian folk remedies to treat headache.

2. Materials and methods

The National Library Service website of the Italian Libraries Network was consulted to find the literature sources. For this purpose, the following key words were entered: “folk medicine”, “usages and customs”, “folk traditions”, “traditional knowledge”, and “folk remedies”. This approach allowed us to unearth about one hundred sources (books and journal articles), written by anthropologists, physicians, ethnographers, folklorists, and scholars of local history.

Among these sources, twenty-seven showed at least one plant-based treatment to heal headache. The plants have been identified as species in two cases: 1) when they were described by their scientific name; 2) when they are mentioned by their Italian name. In the last case, the plant (e.g. onion) has been referred to species (e.g. *Allium cepa* L.) only when there was no doubt. In all other cases, only the genus was attributed (e.g. incense, *Boswellia* spp.).

The names of the plant families were reported according to the guidelines of Angiosperm Phylogeny Group (Stevens, 2001).

Despite new discoveries accumulated over the last 30 years have increased the current knowledge, etiopathogenetic mechanisms of the headache remain unclear and the most common therapeutic approaches are based on analgesic or NSAIDs administration (SISC, 2011). For these reasons, a detailed search of the scientific data banks such as Medline and Scopus was undertaken to uncover recent results concerning the anti-inflammatory, anti-nociceptive, and analgesic activities for each plant-based treatment.

Finally, the collected data were compared with plant-based remedies described in the *Corpus Hippocraticum* (a collection of 62 medical works written between the 5th century BCE and the 2nd century AD that were ascribed to Hippocrates), *Naturalis Historia* (Pliny the Elder, 1st century AD), *De Materia Medica* (Dioscorides, 1st century AD), *Opera Omnia* (Galen, 2nd century AD) and *Liber Medicinalis* (Serenus Sammonicus, 2nd century AD).

3. Results

In the texts of Italian folk medicine, consulted in this work, we have not found different terms to denote different types of cephalalgia. The pain was described generically with the term “headache” or “migraine”, used as synonym. Even if, only in few cases, the consultation of the sources has highlighted types of headache that affects temple, forehead and sometimes extends to the periorbital region. More details, instead, have been claimed about aetiology. Italian folk healers observed that headache could be related to environmental, biological or magical events. Headache was known to be associated to arthritis, cold, blood circulation disorders, oversleeping, fasting and, on the contrary, eating heartily. The exposure to sun, heat, cold and draught as well as sleeping in a field of spring crocus (*Crocus albiflorus* Kit. ex Schult.), meadow saffron (*Colchicum autumnale* L.) and common laburnum (*Laburnum anagyroides* Medik.) was thought to cause headache attacks. Moreover, Italian folk healers believed that magical factors including evil eye, sleeping under the moon or under a walnut tree, watching a toad and combing one's hair at Friday, could induce headache.

The collected data pointed out 74 plants utilized by Italian folk medicine to heal headache belonging to 39 families. Out of these, the most cited were Asteraceae (16.2%) and Lamiaceae (13.5%). This study has highlighted 76 plant-based remedies which were taken through the following routes of administrations: topical (59.2%), oral (27.6%), and inhalation (13.2%). Oral administration was performed swallowing decoctions (38.1), infusions (28.6%), crude plant parts (23.8%), and alcoholic extracts (9.5%). Inhalation was performed breathing steams (50%), sniffing powder (30%), and aspirating juice (20%). The parts of plants used are the following: leaves (27.8%), aerial parts (26.6%), flowers (16.5%), fruits (13.9%), seeds (8.9%), roots (2.5%), tubers (2.5%) and resins (1.3%) (Table 1).

The historical sources consulted in this paper highlighted that, among the 74 plants utilized by Italian folk medicine between the late nineteenth and the early to mid-twentieth century for the treatment of headache, 31 (41.9%) were already included in the pharmacopoeia between the 5th century BCE and the 2nd century AD (Table 1).

Nowadays, a significant percentage (78.4%, N = 58) of plants utilized by Italian folk medicine showed, *in vivo*, *in vitro* or in human trials, an analgesic, anti-nociceptive and anti-inflammatory properties which could explain and justify their potential role in the cure of headache. In Table 2 these pharmacological activities are summarized.

4. Discussion

Natural products contribute significantly to discover new bioactive compounds for development of drugs to treat human diseases. More than half of all approved small-molecule drugs, in fact, show a structure originally ascribable to natural products. In addition, a large number of natural products, nowadays, are used in clinical trials (Stratton et al., 2015).

Similarly, natural treatments, derived from plants and animals, have been an unlimited “reservoir” of remedies since ancient times, and their use has been transmitted for centuries in traditional medicine (Confessor et al., 2009). According to De Vos (2010), natural remedies of the ancient Mediterranean area had a noteworthy lifespan in the European *materia medica*, well established until the 19th century.

In this work, 41.9% of the plant-based treatments utilized by Italian folk medicine between the late nineteenth and the early to mid-twentieth century to heal headache seems to represent the continuity of the medical tradition over about two thousand years.

Several recent studies have shown that knowledge of folk medicine is based upon the pharmacopoeia of ancient Greece and Rome (Pollio et al., 2008; Leonti et al., 2009; Tagarelli et al., 2010, 2013), even if the biggest obstacle is represented by the impossibility to state with certainty which plants (and diseases) are which. These difficulties are not insurmountable. An interdisciplinary approach with philological,

Table 1
Plants used by Italian folk medicine to treat headache and mentioned to serve the same purpose, by historical sources (5th century BCE–2nd century AD).

Family/scientific/common name	Italian name	Plant parts	Route of administration	References	Historical sources	References
Adoxaceae <i>Sambucus nigra</i> L. (Elderberry)	Sambuco	Fr, Lf	Use suffumigations of elderberry flowers boiled in the water; Smear on the temples elderberry leaves	Ostermann (1894); Nardi (1935)	Hippocrates (<i>De Morbis II</i> , 19).	(Potter, 1988a)
Amarantaceae <i>Beta vulgaris</i> L. (Red beet)	Bieta	Ap	Aspirate by nose a juice mixture obtained pressing the following plants: anemone, red beet, cabbage, greater celandine, cyclamen, and onion	(Pignatari, 1894)	Hippocrates (<i>De Morbis II</i> , 12; <i>De Affectionibus Intertribus</i> 48). Pliny (<i>Naturalis Historia XX</i> , 27). Dioscorides (<i>De Materia Medica II</i> , 149).	(Potter, 1988a, 1988b) (Plinio, 1985) (Kühn, 1829)
Amaryllidaceae <i>Allium cepa</i> L. (Onion)	Cipolla	Ft	Aspirate by nose a juice mixture obtained pressing the following plants: anemone, red beet, cabbage, greater celandine, cyclamen, and onion	(Pignatari, 1894)	Dioscorides (<i>De Materia Medica II</i> , 180).	(Kühn, 1829)
Apiaceae <i>Foeniculum vulgare</i> Mill. (Fennel)	Finocchio	Lf	Put on the head a cloth moistened with infusion of the following plants: wormwood, wild thyme, marjoram, thyme, fennel, rosemary, sage, and costmary	(Coronedi-Berti, 1877)	Galen (<i>Opera Omnia XII</i> , 568; <i>XIV</i> , 399).	(Kühn, 1826, 1827)
<i>Pimpinella anisum</i> L. (Anise)	Anice	Sd	Inhale powder obtained from a mixture of pyrethrum, white hellebore, betony, anise, and fumitory seeds	(Finamore, 1894)	Pliny (<i>Naturalis Historia XX</i> , 73). Dioscorides (<i>De Materia Medica III</i> , 58).	(Plinio, 1985) (Kühn, 1829)
Asparagaceae <i>Convallaria majalis</i> L. (Wood lily)	Mughetto	Fr	Drink wine infusion of wood lily	(Pedrotti and Bertoldi, 1930)	–	–
<i>Leopoldia comosa</i> (L.) Parl. (Tassel hyacinth)	Cipollaccio	Tb	Smear on the temples bulbs in halves of tassel hyacinth	(Adriano, 1932)	–	–
Asteraceae <i>Achillea arata</i> L. (Black yarrow)	Millefoglio del calcare	Ap	Drink infusion of black yarrow	(Chioyenda-Bensi, 1955)	–	–
<i>Achillea millefolium</i> L. (Yarrow)	Millefoglio	Fr	Drink infusion of yarrow combined in equal amounts with chamomile	(Chioyenda-Bensi, 1955)	–	–
<i>Achillea erba-rotta</i> All. subsp. moschata (Wulfen) I. Richardson (Iva)	Millefoglio del granito	Ap	Drink infusion of iva	(Chioyenda-Bensi, 1955)	–	–
<i>Arctium lappa</i> L. (Burdock)	Bardana maggiore	Lf	Put on the head fresh leaves of burdock mixed in olive oil	(Pedrotti and Bertoldi, 1930)	–	–
<i>Arnica montana</i> L. (Arnica)	Arnica	Fr	Drink spirit in which arnica flowers were soaked	(Chioyenda-Bensi, 1955)	–	–
<i>Artemisia absinthium</i> L. (Wormwood)	Assenzio	Lf	Put on the head a cloth moistened with infusion of the following plants: wormwood, wild thyme, marjoram, thyme, fennel, rosemary, sage, and costmary; Drink decoction of wormwood	(Coronedi-Berti, 1877; Ostermann, 1894)	Galen (<i>Opera Omnia XII</i> , 524).	(Kühn, 1826)
<i>Artemisia</i> spp. (Artemisia)	Artemisia	Ap, Lf	Apply on the forehead a poultice of artemisia and white horehound mixed with vinegar; Drink decoction of artemisia	(Pasquarelli, 1987; Ostermann, 1894)	–	–
<i>Glebionis coronaria</i> (L.) Cass. ex Spach (Garland)	Crisantemo giallo	Fr	Put on the head garland flowers	(Pitrè, 1896)	–	–
<i>Helianthus annuus</i> L. (Sunflower)	Girasole	Sd	Drink Holy Water infusion of sunflower seeds	(Ostermann, 1894)	–	–
<i>Matricaria chamomilla</i> L. (Chamomile)	Camomilla	Fr	Drink infusion of chamomile alone or combined in equal amounts with yarrow; Put on the head a poultice of chamomile; Apply two red hosts impregnated with chamomile water, when the pain was localized in the periorbital area	(Chioyenda-Bensi, 1955; Pedrotti and Bertoldi, 1930; Latronico, 1935)	–	–
<i>Tanacetum balsamita</i> L. (Costmary)	Erba di Santa Maria	Lf	Put on the head a cloth moistened with infusion of the following plants: wormwood, wild thyme, marjoram, thyme, fennel, rosemary, sage, and costmary	(Coronedi-Berti, 1877)	–	–
<i>Tanacetum cinerariifolium</i> (Trevir.) Sch. Bip. (Pyrethrum)	Piretro	Ap	Inhale powder obtained from a mixture of pyrethrum, white hellebore, betony, anise, and fumitory seeds; Chew stavesacre and pyrethrum	(Finamore, 1894; Pignatari, 1894)	–	–
Brassicaceae <i>Brassica oleracea</i> L. (Cabbage)	Cavolo	Lf	Put on the head fresh leaves of cabbage mixed in warm oil; Aspirate	(Ostermann, 1894; Pignatari, 1894)	Hippocrates (<i>De Morbis II</i> , 19).	(Potter, 1988a) (continued on next page)

Table 1 (continued)

Family/scientific/common name	Italian name	Plant parts	Route of administration	References	Historical sources	References
<i>Lunaria annua</i> L. (Honesty)	Erba argentina	Ap	by nose a juice mixture obtained pressing the following plants: anemone, red beet, cabbage, greater celandine, cyclamen, and onion	1894)	Pliny (<i>Naturalis Historia</i> XX, 33).	(Plinio, 1985)
Burseraceae					Galen (<i>Opera Omnia</i> XIV, 318, 579).	(Kühn, 1827)
<i>Boswellia</i> spp. (Incense)	Incenso	Ap	Smear on the temples aerial parts of honesty	(De Giacomo, 1896)	–	–
Campanulaceae			Use suffumigations of incense boiled in the water	(Sembianti, 1936)	–	–
<i>Campanula glomerata</i> L. (Dane's blood)	Campanula a mazzetti	Fr, Lf	Drink decoction of Dane's blood	(Pedrotti and Bertoldi, 1930)	–	–
Cactaceae					–	–
<i>Opuntia ficus indica</i> (L.) Mill. (Cactus pear)	Fico d'India	Ft	Smear on the temples sliced fruits of cactus pear	(Zanetti, 1892)	–	–
Cannabaceae					–	–
<i>Humulus lupulus</i> L. (Hop)	Luppolo	Fr	Eat hop fruits in barley soup	(Pedrotti and Bertoldi, 1930)	–	–
Caprifoliaceae					–	–
<i>Valeriana officinalis</i> L. (Valerian)	Valeriana	Ap, Fr	Wash head with valerian flowers water; Rub on head with valerian flowers; Rub on forehead the whole plant of valerian	(Ostermann, 1894; Ferraro, 1885)	–	–
Crassulaceae					–	–
<i>Sempervivum tectorum</i> L. (Common houseleek)	Semprevivo dei tetti	Ap	Put on the head common houseleek	(De Nino, 1891)	Dioscorides (<i>De Materia Medica</i> IV, 88).	(Kühn, 1829)
Cucurbitaceae					Galen (<i>Opera Omnia</i> XII, 508).	(Kühn, 1826)
<i>Cucumis sativus</i> L. (Cucumber)	Cetriolo	Ft	Smear on the temples peel of cucumber; Apply on the forehead sliced cucumber	(Adriano, 1932; Lombardi-Satriani, 1970; Marzano, 1890)	Hippocrates (<i>De Morbis</i> II, 12).	(Potter, 1988a)
<i>Cucurbita maxima</i> Duchesne (Pumpkin)	Zucca	Ft	Brought pumpkin in halves like a hat	(Pasquarelli, 1987; Lombardi-Satriani, 1970; Marzano, 1890)	Galen (<i>Opera Omnia</i> XII, 570; XIV, 557).	(Kühn, 1826, 1827)
<i>Echallium elaterium</i> (L.) A. Rich. (Squirting cucumber)	Cocomero asinino	Ft	Aspirate by nose a juice obtained pressing squirting cucumber fruit	(Pitrè, 1896)	Dioscorides (<i>De Materia Medica</i> IV, 155).	(Kühn, 1829)
Cupressaceae					Galen (<i>Opera Omnia</i> XIV, 316).	(Kühn, 1827)
<i>Juniperus communis</i> L. (Juniper)	Ginepro	Ft	Use suffumigations of juniper berries boiled in the water	(Sembianti, 1936)	–	–
Ericaceae					–	–
<i>Vaccinium myrtillus</i> L., <i>Vaccinium vitis idaea</i> L. (Bilberry)	Mirtillo	Ft	Drink spirit in which bilberry fruits were soaked	(Pedrotti and Bertoldi, 1930)	–	–
Fabaceae					–	–
<i>Vicia faba</i> L. (Broad bean)	Fava	Sd	Smear on the temples seeds in halves of broad bean	(Zanetti, 1892)	Galen (<i>Opera Omnia</i> XIV, 399, 548).	(Kühn, 1827)
Gentianaceae					–	–
<i>Gentiana lutea</i> L. (Gentian)	Gentiana	Ap	Drink decoction of gentian	(De Nino, 1891)	–	–
Juglandaceae					–	–
<i>Juglans regia</i> L. (Walnut)	Noce	Lf	Put on the head fresh leaves of walnut	(Pitrè, 1896)	Pliny (<i>Naturalis Historia</i> XXIII, 45).	(Plinio, 1985)
Hypericaceae					Galen (<i>Opera Omnia</i> XII, 569).	(Kühn, 1826)
<i>Hypericum hircinum</i> L. (Stinking tutsan)	Ruta caprina	Ap	Bring on the head a bundle of stinking tutsan	(Geraci, 1957; Pagano, 1992; Lanza, 2006)	–	–
Iridaceae					–	–
<i>Iris germanica</i> L. (German iris)	Giaggiolo	Rt	Spread on the head pounded root of German iris	(Pedrotti and Bertoldi, 1930)	Pliny (<i>Naturalis Historia</i> XXI, 83).	(Plinio, 1985)
Lamiaceae					Dioscorides (<i>De Materia Medica</i> I, 1).	(Kühn, 1829)
<i>Lavandula angustifolia</i> Mill., <i>Lavandula stoechas</i> L. (Lavender)	Lavanda	–	–	(Ostermann, 1894)	Galen (<i>Opera Omnia</i> XII 503, 511, 556, 558, 569, 579, 581, 597; XIV, 316).	(Kühn, 1826, 1827)

(continued on next page)

Table 1 (continued)

Family/scientific/common name	Italian name	Plant parts	Route of administration	References	Historical sources	References
<i>Marrubium vulgare</i> L. (White horehound)	Marrubio	Ap	Apply on the forehead a poultice of artemisia and white horehound mixed with vinegar	(Pasquardelli, 1987)		
<i>Mentha pulegium</i> L. (Pennyroyal)	Puleggio	Ap	Use suffumigations of pennyroyal boiled in the wine	(Pitrè, 1896)		Pliny (<i>Naturalis Historia</i> XX, 54). Galen (<i>Opera Omnia</i> XII, 524, 568; XIV, 399). Serenus Sammonicus (<i>Liber Medicinalis</i> I, 15). Galen (<i>Opera Omnia</i> XII 503, 511, 512). (Kühn, 1826, 1827) (Pépin, 1950) (Kühn, 1826)
<i>Origanum majorana</i> L. (Marjoram)	Maggiorana	Ap, Lf	Put on the head a cloth moistened with infusion of the following plants: wormwood, wild thyme, marjoram, thyme, fennel, rosemary, sage, and costmary; Bring on the head marjoram in a sack; Inhale dried and pulverized leaves of marjoram	(Coronedi-Berti, 1877; Ostermann, 1894; Pitrè, 1896)		
<i>Rosmarinus officinalis</i> L. (Rosemary)	Rosmarino	Lf	Put on the head a cloth moistened with infusion of the following plants: wormwood, wild thyme, marjoram, thyme, fennel, rosemary, sage, and costmary	(Coronedi-Berti, 1877)		
<i>Salvia officinalis</i> L. (Sage)	Salvia	Lf	Put on the head a cloth moistened with infusion of the following plants: wormwood, wild thyme, marjoram, thyme, fennel, rosemary, sage, and costmary; Drink decoction of sage	(Coronedi-Berti, 1877; Chiovenda-Bensi, 1955)		
<i>Stachys officinalis</i> (L.) Trevis. (Betony)	Betonica	Ap	Inhale powder obtained from a mixture of pyrethrum, white hellebore, betony, anise, and fumitory seeds; Spread on the head dried and powdered betony mixed with honey; Drink decoction of betony collected during St. John's night	(Finamore, 1894; Pedrotti and Bertoldi, 1930; Zanetti, 1892)		Dioscorides (<i>De Materia Medica</i> IV, 2). (Kühn, 1829)
<i>Teucrium</i> spp.	Camedrio	–	–	(Tancredi, 1938)		
<i>Thymus serpyllum</i> L. (Wild thyme)	Serpillo	Lf	Put on the head a cloth moistened with infusion of the following plants: wormwood, wild thyme, marjoram, thyme, fennel, rosemary, sage, and costmary	(Coronedi-Berti, 1877)		Pliny (<i>Naturalis Historia</i> XX, 90). Dioscorides (<i>De Materia Medica</i> III, 40). Galen (<i>Opera Omnia</i> XII, 512, 556, 558, 579, 597) (Kühn, 1826)
<i>Thymus</i> spp.	Timo	Lf	Put on the head a cloth moistened with infusion of the following plants: wormwood, wild thyme, marjoram, thyme, fennel, rosemary, sage, and costmary	(Coronedi-Berti, 1877)		
Lauraceae						
<i>Laurus nobilis</i> L. (Bay laurel)	Alloro	Lf	Spread forehead with oil extracted from bay laurel	(Nardi, 1935)		Pliny (<i>Naturalis Historia</i> XXIII, 43, 80). Galen (<i>Opera Omnia</i> XII 503, 512, 514, 556, 558, 570, 579, 580, 581, 597). (Plinio, 1985) (Kühn, 1826)
Melanthiaceae						
<i>Paris quadrifolia</i> L. (Herb Paris)	Uva di volpe	Lf	Put on the head fresh leaves of herb paris	(Ostermann, 1894)		
<i>Veratrum album</i> L. (White hellebore)	Elleboro bianco	Ap	Inhale powder obtained from a mixture of pyrethrum, white hellebore, betony, anise, and fumitory seeds	(Finamore, 1894)		Hippocrates (<i>De Morbis</i> II, 12). Pliny (<i>Naturalis Historia</i> XXV, 89) (Plinio, 1985)
Oleaceae						
<i>Olea europea</i> L. (Olive)	Olivo	Ft	Spread head with olive oil	(Bernoni, 1878)		Pliny (<i>Naturalis Historia</i> XXIII, 34). Galen (<i>Opera Omnia</i> XII, 513, 558, 593, 596; XIV, 315, 316, 318). (Plinio, 1985) (Kühn, 1826, 1827)
Papaveraceae						
<i>Chelidonium majus</i> L. (Greater celandine)	Celidonia	Ap	Aspirate by nose a juice mixture obtained pressing the following plants: anemone, red beet, cabbage, greater celandine, cyclamen, and onion	(Pignatari, 1894)		Serenus Sammonicus (<i>Liber Medicinalis</i> I, 23). (Pépin, 1950)
<i>Fumaria officinalis</i> L. (Fumitory)	Fumaria	Ap	Inhale powder obtained from a mixture of pyrethrum, white hellebore, betony, anise, and fumitory seeds	(Finamore, 1894; Ferraro, 1885)		
Pinaceae						
<i>Abies alba</i> Mill. (Silver fir)	Abete	Rs	Smear on the temples silver fir resin	(Pedrotti and Bertoldi, 1930)		
Plantaginaceae						
<i>Plantago</i> spp. (Plantain)	Piantaggine	Ap	Smear on the temples plantain pounded in vinegar and mixed with alumen	(Ostermann, 1894)		Galen (<i>Opera Omnia</i> XII, 508). (Kühn, 1826)
Poaceae						
<i>Avena sativa</i> L. (Oat)	Avena	Sd	Eat oat seeds put in the water and choosing the floating ones	(Ferraro, 1885)		

(continued on next page)

Table 1 (continued)

Family/scientific/common name	Italian name	Plant parts	Route of administration	References	Historical sources	References
<i>Triticum</i> spp. (Wheat) Poaceace	Frumento, Grano	Sd	Smear on the temples wheat flour mixed with vinegar	(Nardi, 1935)	–	–
<i>Portulaca oleracea</i> L. (Verdolaga)	Portulaca	Lf	Eat verdolaga salad; Apply both on the temples and forehead leaves of verdolaga	(Coronedi-Berti, 1877; Finamore, 1894)	Pliny (<i>Naturalis Historia</i> XX, 81). Dioscorides (<i>De Materia Medica</i> II, 150). Galen (<i>Opera Omnia</i> XII, 508; XIV, 315)	(Plinio, 1985) (Kühn, 1829) (Kühn, 1826, 1827)
Primulaceae						
<i>Cyclamen</i> spp. (Cyclamen)	Ciclamino	Ap, Lf	Put on the head cyclamen mixed with fat and butter; Aspirate by nose a juice mixture obtained pressing the following plants: anemone, red beet, cabbage, greater celandine, cyclamen, and onion	(Ostermann, 1894; Pignatari, 1894)	Pliny (<i>Naturalis Historia</i> XXV, 84). Dioscorides (<i>De Materia Medica</i> II, 193).	(Plinio, 1985) (Kühn, 1829)
<i>Primula veris</i> L. (Cowslip) Ranunculaceae	Primula	Ap	Put on the head a hot pack of cowslip	(Pedrotti and Bertoldi, 1930)	–	–
<i>Anemone</i> spp. (Anemone)	Anemone	Ap	Aspirate by nose a juice mixture obtained pressing the following plants: anemone, red beet, cabbage, greater celandine, cyclamen, and onion	(Pignatari, 1894)	Pliny (<i>Naturalis Historia</i> XXI, 94). Galen (<i>De Materia Medica</i> II, 207).	(Plinio, 1985) (Kühn, 1829)
<i>Delphinium staphysagria</i> L. (Stavesacre) Rubiaceae	Stafisagria	Ap	Chew stavesacre and pyrethrum	(Pignatari, 1894)	–	–
<i>Coffea</i> spp. (Coffee)	Caffè	Sd	Apply on forehead toasted and powdered coffee; Use suffumigations of coffee powder boiled in the water	(Pitrè, 1896; Bernoni, 1878)	–	–
Rhamnaceae						
<i>Paliurus spina-christi</i> Mill. (Christ's thorn) Rutaceae	Spina Christi	Lf	Put on the head pounded leaves of Christ's thorn	(Calvia, 1927)	–	–
<i>Citrus limon</i> (L.) Osbeck (Lemon)	Limone	Ft	Apply on the forehead fresh or roasted sliced lemon	(Pitrè, 1896; Pignatari, 1894; De Nino, 1891; Bernoni, 1878)	–	–
<i>Ruta</i> spp. (Rue)	Ruta	Ap	Bring on the head rue in a sack	(Adriano, 1932; Mancarella, 1930)	Pliny (<i>Naturalis Historia</i> XX, 51). Dioscorides (<i>De Materia Medica</i> III, 45). Galen (<i>Opera Omnia</i> XII, 512, 514, 556, 558, 568, 569, 579, 581, 597; XIV, 500, 516, 543).	(Plinio, 1985) (Kühn, 1829) (Kühn, 1826, 1827)
Santalaceae						
<i>Viscum album</i> L. (Mistletoe)	Vischio	Ap	Drink decoction of mistletoe	(Zanetti, 1892)	Serenus Sammonicus (<i>Liber Medicinalis</i> I, 17)	(Pépin, 1950)
Solanaceae						
<i>Capsicum annuum</i> L. (Pepper)	Pepe	Ft	Apply on forehead powdered pepper mixed with an egg; Chew pepper with raisins	(Pitrè, 1896; Pignatari, 1894; Lombardi-Satriani, 1970)	–	–
<i>Hyoscyamus niger</i> L., <i>Hyoscyamus albus</i> L. (Henbane) Nicotiana glauca L. (Tobacco)	Giusquiamo Tabacco	Lf, Rt Lf	Drink decoction of henbane root; Smear on the temples henbane leaves; put on the head fresh leaves of henbane Inhale snuff tobacco	(Pedrotti and Bertoldi, 1930; Zanetti, 1892; Nardi, 1935) (Pasquarelli, 1987; Ostermann, 1894) (Ferraro, 1885)	–	–
<i>Solanum nigrum</i> L. (Black nightshade)	Solano	Ap	Put on the head black nightshade	(Pitrè, 1896; Pignatari, 1894; Lombardi-Satriani, 1970; Bertagnon, 1955)	Hippocrates (<i>De Morbis</i> III, 1). Pliny (<i>Naturalis Historia</i> XXVII, 44). Galen (<i>Opera Omnia</i> XII, 508, 509; XIV, 516).	(Potter, 1988b) (Plinio, 1985) (Kühn, 1826, 1827)
<i>Solanum tuberosum</i> L. (Potato)	Patata	Tb	Apply on forehead crude or roasted sliced potatoes	(Pitrè, 1896; Adriano, 1932; Lombardi-Satriani, 1970; Bertagnon, 1955)	–	–
Verbenaceae						
<i>Verbena officinalis</i> L. (Vervain)	Verbena	Ap, Lf	Apply both on the temples and forehead leaves of vervain; Fasten forehead with vervain	(Coronedi-Berti, 1877; Finamore, 1894; Nardi, 1935)	Galen (<i>Opera Omnia</i> XII, 575).	(Kühn, 1826)
Vitaceae						

(continued on next page)

Table 1 (continued)

Family/scientific/common name	Italian name	Plant parts	Route of administration	References	Historical sources	References
<i>Vitis vinifera</i> L. (Grapevine)	Vite	Lf	Smear on the temples minced leaves of grapevine; Chew pepper with raisins	(Ostermann, 1894; Lombardi-Satriani, 1970; Pignatari, 1894)	Pliny (<i>Naturalis Historia</i> XXII, 3). Dioscorides (<i>De Materia Medica</i> V, 1). Galen (<i>Opera Omnia</i> XII, 508; XIV, 315).	(Plinio, 1985) (Kühn, 1829) (Kühn, 1826, 1827)

Species names given in bold indicate those plants that nowadays show analgesic, anti-nociceptive, and anti-inflammatory properties.
Plant parts: Ap = Aerial parts; Fr = Flower; Ft = Fruit; Lf = Leaf; Rs = Resin; Rt = Root; Sd = Seed; Tb = Tuber.

botanic, zoological, anthropological and pharmacological points of view, could be the best way to try to identify natural remedies and diseases described in the ancient texts. Aliotta and colleagues (2003) stated that many scholars who have studied the Hippocratic texts generally agree on the identification of the plant-based prescriptions mentioned in the *Corpus Hippocraticum*. Such a result has been shown by Riddle (1987), who states that only 11 of 257 plants in the *Corpus Hippocraticum* cannot be identified with certainty. Moreover, projects such as GALEN, building semantic libraries for medical terminology, allow us to label medical terminology over the centuries (Buenz et al., 2004).

The data collected in this work show that the plant-based remedies utilized by Italian folk medicine for headache were prepared in several ways, usually depending on the plant used. Some of these methods included infusions, decoctions, fomentations, fumigations and macerations. Others included inhalation of powder or the juice of the plant. Treatments were administered by topical, oral and nasal routes. The most used route of administration was topical for which plants were mixed with vinegar, spirit, honey, olive oil, fat, butter, flour and albumen to increase penetration in the skin of the head. Sometimes, the remedies were given using, also, ritualistic objects or reciting prayers and formulas, while sometimes drawing symbols on the body (Romeo et al., 2015). This way represents a further link with the past. According to Karenberg and Leitz (2001), the ancient physicians did not distinguish between magic and medicine. The Egyptian medical papyri, for example, encompass not only what we define rational descriptions and prescriptions, but also magical and religious elements.

A large numbers of experimental studies have shown that plant compounds exerts anti-inflammatory effects through a variety of different mechanisms. Inflammation is the result of the interaction of a variety of inflammatory mediators, and cross mechanisms that exist between various mediators. Inflammatory cytokines and mediators are key components in the inflammation process; thus, inhibition of these targets is an efficient way to indirectly prevent the occurrence and development of inflammation and reduce the resulting damage (Wang et al., 2013).

From a pharmacological point of view, an overview of the recent literature reveals that a significant percentage (78.4%) of the plants, utilized by Italian folk healers in the late nineteenth and the early to mid-twentieth century to treat headache, contains components such as flavonoids, terpenoids and phenylpropanoids, which exhibit a large spectrum of anti-inflammatory, anti-nociceptive and analgesic activities. In fact, the results obtained by recent *in vivo* and *in vitro* studies have demonstrated that secondary metabolites show inhibitory activity against the NF- κ B, NO, COX-2, TNF- α pathways which play an important role in triggering different types of headaches (Gorji, 2003; Capuano et al., 2009; Domingues et al., 2015; Reuter et al., 2002).

NF- κ B is a transcription factor that can be induced by a wide variety of stimuli, including stress, bacteria, viruses, cytokines, and free radicals. It is a ubiquitous protein, composed mainly of two proteins, p50 and p65, that regulates the expression of genes encoding the pro-inflammatory cytokines, chemokines and inducible enzymes. (Baeuerle and Henkel, 1994). Among polyphenols extracted from *Vaccinium myrtillus* fruits, quercetin, epicatechin, and resveratrol were potent inhibitors of NF- κ B activation in a human monocytic cell line (Karlsen et al., 2010). Crude extract of *Ruta graveolens* plant, the diethyl ether fraction and the isolated active compound, Rg 001 (3-(1',1'-dimethylallyl)-6-hydroxy-7-methoxy-chromen-2-one) repress activation of p65/NF- κ B in LPS macrophage cells by inhibiting the activation of I κ B α and effectively suppress nuclear translocation of NF- κ B (Raghav et al., 2007). Finally, bioassay guided fractionation of the *Valeriana officinalis* extracts led to the isolation of acetylvalerenolic acid and valerenic acid, two sesquiterpenes active as inhibitors of NF- κ B (Jacobo-Herrera et al., 2006).

NO is an important inflammatory mediator acting on vascular vasodilation and increasing permeability to facilitate the infiltration and

Table 2
Plants used by Italian folk medicine to treat headache, between the late nineteenth century and the early to mid-twentieth century, and their pharmacological properties at the present time.

Family/scientific/ common name	Part of plant	Type of extract	Components	Experimental models	Animal or cell	Dose range	Minimal active concentration	Anti- inflammatory, analgesic pathways	Positive Controls	References
<i>Adoxaceae</i> <i>Sambucus nigra</i> L. (Elderberry)	Fruit	Methanolic fraction	TP3	AA Enzyme-linked immunosorbent assay	-	10 µg/ml	-	COX-2	Indomethacin	Thole et al., 2006
	Fruit	Acetone extract	TP4	AA Enzyme-linked immunosorbent assay	-	10 µg/ml	-	COX-2	Indomethacin	
<i>Amarandaceae</i> <i>Beta vulgaris</i> L. (Red beet)	Root	Ethanol extract	-	Xylene-induced ear edema	Mice	200–400 mg/kg	400 mg/kg	-	-	Atta and Alkofahi, 1998
	Root	Ethanol extract	-	Cotton pellet granuloma	Rats	200–400 mg/kg	400 mg/kg	-	-	
<i>Amaryllidaceae</i> <i>Allium cepa</i> L. (Onion)	Bulb	-	ALC-02	Carrageenan-induced paw edema	Rats	50–200 mg	200 mg	Anti-histaminic mediators	Ibuprofen	Kaiser et al., 2009
	Bulb	Fresh onion Juice	-	Carrageenan-induced paw edema	Rats	5–10 ml/kg	7.5 ml/kg	Cox	Diclofenac; morphine	Nasri et al., 2012
	Bulb	Fresh onion Juice	-	Hot Plate	Mice	5–10 ml/kg	7.5 ml/kg	-	Morphine	
	Bulb	Fresh onion Juice	-	Formalin test	Rats	-	7.5 ml/kg	Cox	Morphine	
<i>Apiaceae</i> <i>Foeniculum vulgare</i> Mill. (Fennel)	Fruit	Methanolic extract	-	Carrageenan-induced paw edema	Mice	-	200 mg/kg	COX; LOX	Indomethacin	Choi and Hwang, 2004
	Fruit	Methanolic extract	-	Arachidonic acid- induced ear edema	Mice	-	200 mg/kg	-	Indomethacin	
	Fruit	Methanolic extract	-	Formaldehyde-induced arthritis	Mice	-	200 mg/kg	-	Indomethacin	
<i>Pimpinella anisum</i> L. (Anise)	Fruit	Methanolic extract	-	Hot plate	Mice	-	200 mg/kg	-	-	
	Fruit	Ethanol extract	-	LPS-stimulated	RAW 264.7 cells	50–250 µg/ml	72.7 µg/ml	NO	Indomethacin	Conforti et al., 2010
	Fruit	-	Trans-anethole	LPS-induced NO production	RAW 264.7 cells	50–250 µg/ml	102.7 µg/ml	NO	Indomethacin	
	Fruit	-	Limonene	LPS-induced NO production	RAW 264.7 cells	50–250 µg/ml	70.1 µg/ml	NO	Indomethacin	
<i>Asteraceae</i> <i>Achillea millefolium</i> L. (Yarrow)	Aerial parts Flower heads	-	Chamazulene carboxylic acid	Ear lobe edema	Mice	50 µM	-	COX-2	Nimesulide	Ramadan et al., 2006
	Aerial parts Flower heads	-	Chamazulene carboxylic acid	Carrageenan-induced paw edema	Rats	300 mg/kg	-	COX-2	Aspirin	
	Aerial parts Flower heads	-	Sesquiterpene Lactones	Respiratory burst assay	Human neutrophils	-	500 µg/ml	-	Indomethacin	Choudhary et al., 2007
<i>Arctium lappa</i> L. (Burdock)	Seeds	Lignans	Lappaol F	LPS-induced NO production	RAW 264.7 cells	-	9.5 µM	NO	Aminoguanidine	Park et al., 2007
	Seeds	Lignans	Diarctigenin	LPS-induced NO production	RAW 264.7 cells	-	9.6 µM	NO	Aminoguanidine	
	Seeds	Lignans	Diarctigenin	LPS-induced NO production	RAW 264.7 cells	6–12 µM	8 µM	NO, PGE, IL, TNF-α	-	Kim et al., 2008
	Seeds	Lignans	Diarctigenin	Zimosan-induced NO production	RAW 264.7 cells	6–12 µM	9.6 µM	NO, PGE, IL, TNF-α	-	
<i>Arnica montana</i> L. (Arnica)	Flowers	Alcoholic extract	-	Carrageenan-induced paw edema	Rats	-	0.1 ml	-	-	Maçêdo et al., 2004
	Flowers	Alcoholic extract	-	Nystatin-induced paw edema	Rats	-	0.1 ml	-	-	

(continued on next page)

Table 2 (continued)

Family/scientific/ common name	Part of plant	Type of extract	Components	Experimental models	Animal or cell	Dose range	Minimal active concentration	Anti- inflammatory, analgesic pathways	Positive Controls	References
<i>Artemisia absinthium</i> L. (Wormwood)	Flowers	Alcoholic extract	-	Histamina induced- vascular permeability LPS-stimulated	Rats	-	0.1 ml	-	-	Lee et al., 2004
<i>Artemisia</i> spp. (Artemisia)	Plant	-	Flavonoid p7F		RAW 264.7	50–200 µg/ml	200 µg/ml	COX-2; PGE2; NO; NF-κB	Naringenin	
	Aereal parts	Essential oil	-	Formalin-induced hind- paw licking	Rats	10–300 mg/kg	100 mg/kg	-	Morphine	Maham et al., 2013
	Aereal parts	Essential oil	-	Hot plate test	Rats	10–300 mg/kg	100 mg/kg	-	Morphine	
	Aereal parts	Essential oil	-	Acetic acid- induced writhing test	Mice	10–300 mg/kg	10 mg/kg	PGE	Morphine; Naloxone	
	Aereal parts	Ethanol extract	p-hydroxyacetophenone	Acetic acid- induced writhing test	Mice	20–500 mg/kg	100 mg/kg	TNF-α; IL-6; COX-2; NO	Indomethacin	Chou et al., 2012
	Aereal parts	Ethanol extract	p-hydroxyacetophenone	Formalin-induced -paw licking	Mice	20–500 mg/kg	100 mg/kg	TNF-α; IL-6; COX-2; NO	Indomethacin	
	Aereal parts	Ethanol extract	p-hydroxyacetophenone	Carrageenan-induced paw edema	Rats	20–500 mg/kg	100 mg/kg	TNF-α; IL-6; COX-2; NO	Indomethacin	
<i>Glebionis coronaria</i> (L.) Cass. ex Spach (Garland)	-	Lyophilized ethanolic extract	-	LPS-induced cells	P388D1 Murine monocyte	-	-	NO; TNF-α	-	Strzelecha et al., 2005
<i>Helianthus annuus</i> L. (Sunflower)	Flower head	-	Diterpene acids	LPS-induced cells	RAW 264.7 Murine macrophages	1–20 µM	1 µM	COX-2; NOS-2	-	Diaz-Viciedo et al., 2008
<i>Maricaria chamomilla</i> L. (Chamomile)	Flower head	-	Diterpene acids	TPA mouse-ear edema	Mice	0.25–1 mg/ear	0.25 mg/ear	-	Indomethacin	Drummond et al., 2012
<i>Tanacetum balsamita</i> L. (Costmary)	Flower	Water extract	Apigenin	LPS-stimulated	THP1 macrophages	4–20 µM	4 µM	TNF-α; IL-6	-	
Brassicaceae <i>Brassica oleracea</i> L. (Cabbage)	Aerial parts	Diethyl ether extract	-	Carrageenan-induced paw edema	Rats	25–100 mg/kg	25 mg/kg	-	Indomethacin	Karaca et al., 2009
Burseraceae <i>Boswellia</i> spp. (Incense)	Aerial parts	Glucoside	Indol-3-carbinol	LPS-induced	RAW 264.7	12.5–100 µM	25 µM	NO; IL-6; IL-8	-	Jiang et al., 2013
	Bark	Methanol extract	-	Hot plate test	Mice	200–400 mg/kg	400 mg/kg	-	Indomethacin	Mothana, 2011
	Bark	Methanol extract	-	Acetate acid-induced writhing test	Mice	200–400 mg/kg	400 mg/kg	-	Aspirin	
	Bark	Methanol extract	-	Cotton induced granuloma test	Rats	200–400 mg/kg	400 mg/kg	-	Indomethacin	
	Bark	Methanol extract	-	Carrageenan-induced paw edema	Rats	200–400 mg/kg	400 mg/kg	-	Indomethacin	
Cactaceae <i>Opuntia ficus indica</i> (L.) Mill. (Cactus pear)	Cladodes	Lyophilised extracts	-	IL-1β-induced	Human chondrocytes	200 µg/ml	-	NO; PGE-2	-	Panico et al., 2007
	Cladodes	Ethanol extract	B-Sitosterol	Adjuvant-induced pouch granuloma	Mice	14 mg/kg	-	-	Hydrocortisone	Park et al., 2001
Cannabaceae <i>Humulus lupulus</i> L. (Hop)	Hop pellet	-	Xanthohumol	LPS-induced	BV2 Mouse	0.5–5 µg/ml	2.5 µg/ml	NO; IL; TNF-α	-	Lee et al., 2011

(continued on next page)

Table 2 (continued)

Family/scientific/ common name	Part of plant	Type of extract	Components	Experimental models	Animal or cell	Dose range	Minimal active concentration	Anti- inflammatory, analgesic pathways	Positive Controls	References
Caprifoliaceae <i>Valeriana officinalis</i> L. (Valerian)	Root	Ethyl acetate extract	Valerenic acid	IL-6/Luc Assay	microglial cells HeLa cells	-	100 µg/ml	NF-kB	-	Jacobo-Herrera et al., 2006
Grassulaceae <i>Sempervivum</i> <i>tectorum</i> L. (Common houseleek)	Leaves	Juice	Flavonolglycosides	Acetic Acid writhing test	Mice	720–2880 mg/kg	1440 mg/kg	-	-	Alberti et al., 2012
Cucurbitaceae <i>Cucumis sativus</i> L. (Cucumber)	Fresh fruit	Water extract	-	Tail immersion test	Mice	250–500 mg/kg	500 mg/kg	-	Diclofenac	Kumar et al., 2010
	Fresh fruit	Water extract	-	Acetic Acid writhing test	Mice	250–500 mg/kg	500 mg/kg	-	Diclofenac	
<i>Ecballium elaterium</i> (L.) A. Rich. (Squirting cucumber)	Fruit juice		Cucurbitacin B	Whittle method	Mice	50–400 mg/kg	200 mg/kg	-	Acetylsalicylic Acid	Yesilada et al., 1988
Cupressaceae <i>Juniperus communis</i> L. (Juniper)	Stem/fruits/ leaves	Methanol/water extract	-	Carrageenan-induced paw edema	Mice	-	100 mg/kg	PGE-2	Indomethacin	Akkol et al., 2009
	Stem/fruits/ leaves	Methanol/water extract	-	PGE2-induced hind paw edema	Mice	-	100 mg/kg	PGE-2	Indomethacin	
	Stem/fruits/ leaves	Methanol/water extract	-	p-benzoquinone- induced writhing test	Mice	-	100 mg/kg	PGE-2	Acetylsalicylic acid	
	Stem/fruits/ leaves	Methanol/water extract	-	Hot plate test	Mice	-	100 mg/kg	PGE-2	Morphin	
Ericaceae <i>Vaccinium myrtillus</i> L., <i>Vaccinium</i> <i>vitis idaea</i> L. (Bilberry)	Fruit	Juice	-	Randomized controlled trial	Human volunteers	330 ml/day	-	IL-1 TNF-α	-	Karlsen et al., 2010
	Fruit	-	Polyphenol	LPS-induced cells	U937 Human monocyte cell line	1.50µmol/l	25µmol/l	NF-kB	-	
Gentianaceae <i>Gentiana lutea</i> L. (Gentian)	Roots	Methanol extract	-	Swimming test	Mice	250–500 mg/kg	250 mg/kg	-	-	Öztürk et al., 2002
	Roots	Methanol extract	-	Tail-clip test	Mice	250–500 mg/kg	250 mg/kg	-	-	
	Roots	Methanol extract	-	Tail-immersion test	Mice	250–500 mg/kg	250 mg/kg	-	-	
Jungladiaceae <i>Juglans regia</i> L. (Walnut)	Leaves	Ethanol/ acquaouextract	-	p-benzoquinone- induced writhing test	Mice	-	500 mg/ml	-	Acetylsalicylic acid	Erdemoglu et al., 2003
		Ethanolic extract	-	Carrageenan-induced paw edema	Mice	-	500 mg/kg	-	Indomethacin	
Hypericaceae <i>Hypericum hircinum</i> L. (Stinking tutsan)	Leaves	Methanolic extract	Quercetin	Forced swimming test	Mice	1–100 mg/kg	100 mg/kg	MAO	-	Chimenti et al., 2006
Iridaceae	Leaves	Methanolic extract	Quercetin	-	-	-	0.010 µM	MAO	Toloxatone	(continued on next page)

Table 2 (continued)

Family/scientific/ common name	Part of plant	Type of extract	Components	Experimental models	Animal or cell	Dose range	Minimal active concentration	Anti- inflammatory, analgesic pathways	Positive Controls	References
<i>Iris germanica</i> L. (German iris)	Rhizomes	Methanolic extract		Formalin-induced paw edema	Rats	50–100 mg/kg	50 mg/kg		Dexamethasone	Ibrahim et al., 2012
	Rhizomes	Methanolic extract	Flavonoids	Formalin-induced paw edema	Rats	10 mg/kg	10 mg/kg		Dexamethasone	
Lamiaceae <i>Lavandula angustifolia</i> Mill., <i>Lavandula stoechas</i> L. (Lavender)	Leaves	Essential oil	-	Lavender essential oil inhalation	Human volunteers	15 min inhalation	-	-	-	Sasanejad et al., 2012
<i>Marrubium vulgare</i> L.	Aerial parts	-	Marrubiin	Microvascular leakage test	Mice	1–100 mg/kg	3 mg/kg	-	Diclofenac	Stulzer et al., 2006
(White horehound) <i>Mentha pulegium</i> L. (Pennyroyal)	Aerial parts	Hydro-ethanol extract	-	Carrageenan-induced ear edema	Mice	300 µl/cm ²	-	-	-	Moussaid et al., 2011
<i>Origanum majorana</i> L. (Marjoram)	Aerial part	Essential oil	Carvacrol	Carrageenan-induced mechanical hypernociception	Mice	25–100 mg/kg	50 mg/kg	TNF-α; NO ₂ PGE2	Indomethacin	Guimaraes et al., 2012
	Aerial part	Essential oil	Carvacrol	Carrageenan-induced paw edema	Mice	25–100 mg/kg	25 mg/kg	TNF-α; NO ₂ PGE2	Dexamethasone	
	Aerial part	Essential oil	Carvacrol	TNF-α-induced mechanical hypernociception	Mice	25–100 mg/kg	50 mg/kg	TNF-α; NO ₂ PGE2	Indomethacin	
	Aerial part	Essential oil	Carvacrol	LPS-induced	Murine macrophages	1–100 µg/ml	1 µg/ml	TNF-α; NO ₂ PGE2	-	
<i>Rosmarinus officinalis</i> L. (Rosemary)	Aerial parts	Hexane and ethyl acetate fractions	Carnosol/Betulinic acid	Carrageenan-induced murine pleurisy	Mice	-	2.5 mg/kg	NO; IL-1β; TNF-α	Dexamethasone Indomethacin	Benincá et al., 2011
	Aerial parts	Ethyl acetate fractions	Ursolic acid	Carrageenan-induced murine pleurisy	Mice	-	25 mg/kg	NO; IL-1β; TNF-α	Dexamethasone Indomethacin	
<i>Salvia officinalis</i> L. (Sage)	Leaves	Hydroalcoholic extract	-	Acetic acid writhing test	Mice	10–100 mg/kg	10 mg/kg	-	Ketoprofen	Rodrigues et al., 2012
	Leaves	Hydroalcoholic extract	-	Formalin test	Mice	3–100 mg/kg	3 mg/kg	-	Morphine Naloxone	
	Leaves	Hydroalcoholic extract	-	Glutamate-induced hind paw edema	Mice	3–100 mg/kg	3 mg/kg	-	-	
	Leaves	Hydroalcoholic extract	-	Capsaicin-induced paw edema	Mice	3–100 mg/kg	10 mg/kg	-	Ruthenium red	
	Leaves	Hydroalcoholic extract	-	Cinnamaldehyde- induced paw edema	Mice	3–100 mg/kg	10 mg/kg	-	Camphor	
<i>Stachys officinalis</i> (L.) Trevis. (Betony)	Aerial parts	Water extract	Iridoids	Carrageenan-induced paw edema	Rats	5 mg/kg	-	-	Diclofenac-Na	Háznagy- Radnai et al., 2012
<i>Thymus serpyllum</i> L. (Wild thyme)	-	-	Carvacrol	Carrageenan-induced paw mechanical hypernociception and edema	Mice	25–100 mg/kg	25 mg/kg	TNF-α NO	Dexamethasone	Guimaraes et al., 2012
<i>Thymus</i> spp.	-	-	Carvacrol	TNF-α, dopamine, or PGE-2 induced mouse paw hypernociception	Mice	25–100 mg/kg	50 mg/kg	TNF-α NO	Indomethacin	
	-	-	Carvacrol	Carrageenan-induced	Mice	25–100 mg/kg	25 mg/kg	TNF-α NO	Indomethacin	(continued on next page)

Table 2 (continued)

Family/scientific/ common name	Part of plant	Type of extract	Components	Experimental models	Animal or cell	Dose range	Minimal active concentration	Anti- inflammatory, analgesic pathways	Positive Controls	References
Lauraceae <i>Laurus nobilis</i> L. (Bay laurel)	-	-	Carvacrol	pleurisy LPS-induced Nitrite production	Murine macrophages	1–100 µg/ml	1 µg/ml	TNF-α NO	-	
	Leaves	Essential oil	-	Tail-flick test	Mice	0.015–0.06 ml/kg	0.03 ml/kg	-	Morphine	Sayyah et al., 2003
	Leaves	Essential oil	-	Formalin-induced hind paw edema	Rats	0.125–0.25 ml/kg	0.25 ml/kg	-	Morphine	
	Leaves	Essential oil	-	Formaldehyde-induced hind paw edema	Rats	0.05–0.2 ml/kg	0.05 ml/kg	-	Piroxicam	
	Aerial parts	Water extract	-	LPS induced	BV2 murine macrophage cell line	0.02–0.6 µl/ml	0.02 µl/ml	COX-2; PGE2	Resveratrol	Orlando et al., 2010
Melanthiaceae <i>Veratrum album</i> L. (White hellebore)	-	-	Cis-resveratrol	LPS-induced cells	THP-1 macrophages	1–100 µM	1 µM	IL-1 COX-2	-	Huang et al., 2014
Oleaceae <i>Olea europea</i> L. (Olive)	Fruit	Olive Vegetation Water	-	LPS-treated	BALB7c Mice	5–125 mg/kg	35 mg/kg	TNF-α	-	Bitler et al., 2005
	Fruit	Olive Vegetation Water	-	LPS-induced	THP-1 monocytes	-	0.5 g/L	TNF-α	Dexamethasone	
Papaveraceae <i>Chelidonium majus</i> L. (Greater celandine)	Aerial parts	Methanol extract	-	Collagen-induced arthritis	Mice	40–400 mg/kg	400 mg/kg	TNF-α; IL-6	-	Lee et al., 2007
<i>Pumaria officinalis</i> L. (Fumitory)	Aerial Part	Ethanol extract	-	Carrageenan-induced paw edema	Rats	100–400 mg/kg	100 mg/kg	-	Phenylbutazone	Rao et al., 2007
	Aerial Part	Ethanol extract	-	Hystamin induced hind paw edema	Rats	100–400 mg/kg	100 mg/kg	-	Phenylbutazone	
	Aerial Part	Ethanol extract	-	Cotton pellet induced granuloma	Rats	100–400 mg/kg	100 mg/kg	-	Phenylbutazone	
	Aerial Part	Ethanol extract	-	Hot plate	Mice	100–400 mg/kg	200 mg/kg	-	Phenylbutazone	
	Aerial Part	Ethanol extract	-	Acetic acid writhing test	Mice	100–400 mg/kg	100 mg/kg	-	Acetylsalicylic acid	
Plantaginaceae <i>Plantago</i> spp. (Plantain)	Leaves/seeds	Methanol extract	-	Acetic acid writhing test	Mice	200–400 mg/kg	400 mg/kg	-	Dipyrone	Atta and Abo El-Sooud, 2004
	Leaves	Methanol extract	-	Tail-flick test	Mice	200–400 mg/kg	400 mg/kg	-	Dipyrone	
	Aerial parts	Hydromethanolic extract	-	LC-MS/MS	Human platelets	1–9 mg/ml	0.65 mg/ml	COX-1 12-LOX	Acetylsalicylic acid; Quercetin	Beara et al., 2010
Poaceae <i>Avena sativa</i> L. (Oat)	Whole grain	-	-	NF-κB inhibitory assay cells	Human 293 T cells	-	2 mg/ml	NF-κB	-	Chu et al., 2013
<i>Triticum</i> spp. (Wheat)	Whole meal flour	Hydrophilic extract	Phenolic acid	LPS-stimulated cells	HT-29 Human intestinal cells	8.2–66 µg/ml	66 µg/ml	IL-8	-	Laddomada et al., 2015
	Whole meal flour	Lipophilic extract	Isoprenoids	LPS-stimulated cells	HT-29 Human intestinal cells	0.01–0.2 µg/ml	0.2 µg/ml	IL-8	-	
Portulacaceae <i>Portulaca oleracea</i> L. (Verdolaga)	Aerial parts	Ethanol extract	-	Hot-plate test	Mice	200–400 mg/kg	400 mg/kg	-	Diclofenac	Chan et al., 2000

(continued on next page)

Table 2 (continued)

Family/scientific/ common name	Part of plant	Type of extract	Components	Experimental models	Animal or cell	Dose range	Minimal active concentration	Anti- inflammatory, analgesic pathways	Positive Controls	References
Primulaceae <i>Cyclamen</i> spp. (Cyclamen)	Aerial parts	Ethanol extract	-	Tail-flick test	Rats	200–400 mg/kg	400 mg/kg	-	Diclofenac	Speroni et al., 2007
	Aerial parts	Ethanol extract	-	Carrageenan-induced paw edema	Rats	200–400 mg/kg	200 mg/kg	-	Diclofenac	
	Aerial parts	Ethanol extract	-	Cotton pellet induced granuloma	Rats	200–400 mg/kg	400 mg/kg	-	Diclofenac	
Ranunculaceae <i>Anemone</i> spp. (Anemone)	Dried Tubers	Petroleum ether/ methanol/chloroform extracts	-	Carrageenan-induced paw edema	Rats	75–150 mg/kg	75 mg/kg	-	Indomethacin	Sun et al., 2011
	Dried Tubers	Petroleum ether/ methanol/chloroform extracts	-	Acetic acid writhing test	Mice	50–100 mg/kg	150 mg/kg	-	Morphin	
Rubiaceae <i>Coffea</i> spp. (Coffe)	Rhizome	Saponins	Radddeanosiide R1, Radddeanosiide F, Radddeanosiide H	Acid acetic-induced pain reaction	Mice	-	125 mg/kg	-	-	Muhammad et al., 2008
	Unroasted beans	Diterpenes	Cafestol; Kahweol	Cell based assay for inhibition of COX-2 activity	RAW 264.7	0.25–5.0 µg/ml	-	COX-2	-	
Rutaceae <i>Citrus limon</i> (L.) Osbeck (Lemon)	Leaves	Essential oil	Mono terpenes	Acetic acid-induced writhing	Mice	50–150 mg/kg	50 mg/kg	-	Morphine	Campelo et al., 2011
	Leaves	Essential oil	Mono terpenes	Formalin test	Mice	50–150 mg/kg	100 mg/kg	-	Morphine; Aspirin	
	Leaves	Essential oil	Mono terpenes	Hot plate test	Mice	50–150 mg/kg	-	-	Morphine	
	Aerial parts	Polyphenolic/ Alkaloid fraction	-	Carrageenan-induced paw edema	Rats	5–15 mg/kg	5 mg/kg	COX-2; 5-LOX	Diclofenac	
	Aerial parts	PhenolicAlkaloid fraction	-	Adjuvant-induced hind paw arthritis	Rats	10 mg/kg	-	COX-2; 5-LOX	Indomethacin	
Santalaceae <i>Viscum album</i> L. (Mistletoe)	Whole plant	Methanolic extract	-	LPS-stimulated	J774A.1 murine macrophage cells	300–500 µg/ml	300 µg/ml	TNF-α; IL; NF-κB	L-NAME	Raghav et al., 2007
	Whole plant	Methanolic extract	3-(1'-1'-dimethyl- allyl)-6-hydroxy-7- methoxy-coumarin	LPS-induced	BALB/c mice	1 mg/25 g	-	NO; IL	-	
	Leaves and stems	Ethyl acetate fraction	Isoflavonoids	P-benzoquinone- induced writhing test	Mice	125–250 mg/kg	125 mg/kg	-	Acetylsalicylic acid	
Solanaceae <i>Capsicum annuum</i> L. (Pepper)	Leaves and stems	Ethyl acetate fraction	Isoflavonoids	Carrageenan-induced hind paw edema	Mice	125–250 mg/kg	125 mg/kg	-	Indomethacin	Orhan et al., 2006
	Fruit	Petroleum ether extract	Carotenoids	Acetic acid-induced writhing	Mice	5–80 mg/kg	5 mg/kg	-	Ibuprofen	
	Fruit	Petroleum ether extract	Carotenoids	Hot plate test	Mice	5–80 mg/kg	80 mg/kg	-	Indomethacin	
<i>Hyoscyamus niger</i> L., <i>Hyoscyamus</i> <i>albus</i> L. (Henbane)	Fruit	Petroleum ether extract	Carotenoids	Carrageenan-induced paw edema	Mice	5–80 mg/kg	5 mg/kg	-	Indomethacin	Hernandez- Ortega et al., 2012
	Seeds	Methanolic extract	-	Hot plate test	Mice	100–400 mg/kg	100 mg/kg	-	Pentazocine	
	Seeds	Methanolic extract	-	Acetic Acid writhing	Mice	100–400 mg/kg	200 mg/kg	-	Acetylsalicylic acid	

(continued on next page)

Table 2 (continued)

Family/scientific/ common name	Part of plant	Type of extract	Components	Experimental models	Animal or cell	Dose range	Minimal active concentration	Anti- inflammatory, analgesic pathways	Positive Controls	References
<i>Solanum nigrum</i> L. (Black nightshade)	Seeds	Methanolic extract	-	test						
				Carrageenan-induced paw edema	Rats	50–200 mg/kg	50 mg/kg	-	Phenibutazone	
	Seeds	Methanolic extract	-	Cotton pellet induced granuloma	Rats	100–400 mg/kg	100 mg/kg	-	Indomethacin	
	Leaves	Chloroform extract	-	Acetic Acid writhing test	Mice	20–200 mg/kg	20 mg/kg	-	Acetylsalicylic acid	Zakaria et al., 2006
	Leaves	Chloroform extract	-	Hot plate-test	Mice	20–200 mg/kg	20 mg/kg	-	Morphine	
	Leaves	Chloroform extract	-	Formalin-test	Rats	20–200 mg/kg	20 mg/kg	-	Acetylsalicylic acid	
<i>Solanum tuberosum</i> L. (Potato)	Leaves	Chloroform extract	-	Carrageenan-induced paw edema	Rats	100–200 mg/kg	100 mg/kg	-	Acetylsalicylic acid	
	Tuber	Ethanol extract	-	Formalin-induced paw licking test	Mice	100–200 mg/kg	100 mg/kg	-	Aminopyrine	Choi and Koo, 2005
	Tuber	Ethanol extract	-	Acetic Acid writhing test	Mice	100–200 mg/kg	100 mg/kg	-	Aminopyrine	
	Tuber	Ethanol extract	-	Hot plate-test	Mice	100–200 mg/kg	100 mg/kg	-	Aminopyrine	
	Tuber	Ethanol extract	-	Carrageenan-induced paw edema	Mice	100–200 mg/kg	100 mg/kg	-	Indomethacin	
	Tuber	Ethanol extract	-	Arachidonic acid- induced ear edema	Mice	100–200 mg/kg	100 mg/kg	-	Indomethacin	
Verbenaceae <i>Verbena officinalis</i> L. (Vervain)	Leaves	Methanolic extract	-	Carrageenan-induced paw edema	Rats	1–3% (50% Methanolic extract)	1%	-	Piroxicam	Calvo, 2006
	Leaves	Methanolic extract	-	Formalin test	Rats	1–3% (50% Methanolic extract)	2.5%	-	Methyl salicylate	
	Leaves	Water extract	-	Carrageenan-induced hind paw edema	Mice	100–400 mg/kg	100 mg/kg	-	Indomethacin	Kosar et al., 2007
	Leaves	Water extract	-	P-benzoquinone- induced writhing test	Mice	100/400 mg/kg	100 mg/kg	-	Acetylsalicylic acid	

penetration of inflammatory mediators, pain substances into the inflammatory site (Wang et al., 2013). Two known lignans, lappacol F, and diartigenin, isolated from *Arctium lappa* seeds, strongly inhibited NO production in the LPS-stimulated RAW264.7 cells (Park et al., 2007).

TNF- α is one of the major pro-inflammatory cytokines that activates monocyte-macrophage cells to release large amounts of IL-1, IL-6, IL-8, PGE₂ and other inflammatory mediators and stimulate a chain reaction of inflammation (Wittmann et al., 1996).

Indol-3-carbinol (I3C) is an autolysis product of glucosinolate present in Brassica plants. I3C reduced the amounts of IL-6 and TNF- α in broncho-alveolar lavage fluid and in the acute lung injury in the LPS-induced mouse model (Jiang et al., 2013). Carnosol, betulinic acid and ursolic acid, isolated from crude extract of *Rosmarinus officinalis* aerial parts have been showed an important anti-inflammatory activity by inhibition not only of leukocytes and exudation, but also of a pro-inflammatory enzyme and mediators (NOx, IL-1b, and TNF- α) in carrageenan-induced murine pleurisy (Benincá et al., 2011). p7F (5,6,3',5'-tetramethoxy 7,4'-hydroxyflavone), a flavonoid extracted from the aerial parts of *Artemisia absinthium*, markedly inhibited PGE₂ activity in LPS-induced macrophages, as shown by Lee and colleagues (2004).

COX-2 belongs to the cyclooxygenase family which are key enzymes of the arachidonic acid cascade (Ramadan et al., 2006). In resting cells, it is not expressed; however, after stimulation by an inflammatory response, COX-2 can be synthesized quickly and is involved in inflammatory processes (Wang et al., 2013).

Three diterpene acids: grandiflorolic, kaurenoic and trachylobanoic acids extracted from *Helianthus annuus* flower head showed inhibition of the expression of COX-2 and the release of inflammatory cytokines in LPS-activated RAW 264.7 macrophages (Díaz-Viciedo et al., 2008).

In addition, MAO inhibitors, usually, are used in the cure of depression and other psychological disorders. They increase the concentration of neuroamines, especially 5-HT and NE. Vasodilatation, which is a source of pain in headache (Shevel, 2011), is inhibited by the depletion of the aforesaid neuroamines. For these reasons, the use of MAO inhibitors is reserved for chronic daily headache patients whose symptoms have not been relieved by conventional prophylactic treatment (Redillas and Solomon, 2000). Extracts from leaves of *Hypericum hircinum* showed MAO-A inhibitory activities. Quercetin showed a selective MAO-A inhibitory activity in the forced swimming test (Chimenti et al., 2006).

Moreover, essential oil from lavender has been studied in a placebo-controlled clinical trial. In this study, the percentage of responders was significantly higher in the lavender cohort than in the placebo cohort, so the inhalation of lavender essential oil may be an efficacious and innocuous route of administration in acute management of migraine headache and his associated symptoms such as photophobia, phonophobia, nausea and vomiting (Sasannejad et al., 2012).

Finally, metabolites isolated from anemone, arnica, red beet, incense, pepper, lemon, cucumber, cyclamen, squirting cucumber, fumitory, gentian, henbane, German iris, walnut, marjoram, pennyroyal, verdolaga, sage, common houseleek, black nightshade, potato, betony, costmary, vervain, mistletoe and grapevine have showed a more generic central and/or peripheral depressant action whose mechanisms are not readily apparent.

Although most, if not all of the plant-based compounds are still in the pre-clinical phase, we speculate that Italian folk medicine exerting anti-headache effect may mainly act through the regulation of Nitric Oxide, Histamine, Serotonin and Arachidonic Acid pathways. Besides, as a holistic medicine, bioactive compounds, isolated from the plants utilized by Italian healers between the late nineteenth century and the early to mid-twentieth century, could be helpful not only to cure headache, but also for the treatment of some other diseases which exhibit an inflammatory component such as cancer, rheumatoid arthritis, sepsis, atherosclerosis and Alzheimer's disease (Calixto et al., 2003, 2004).

5. Conclusions

Folk medicine, based on centuries of experience, is like an uncontrolled laboratory which, if examined carefully and critically, would help us to gain valuable information to be exploited in modern medicine.

Besides historical prospective, this study unearths interesting remedies that can provide a comprehensive basis to perform further investigations for the development of effective drugs against headache.

Even if traditional knowledge about natural remedies to treat several diseases was handed down almost entirely orally, it has been the subject of historical documents, too. These valuable sources, nowadays, should be explored systematically.

The award of the 2015 Nobel Prize for Physiology or Medicine to Prof. Youyou Tu for her discovery and development of artemisinin that is the most effective drug against malaria, gives us a lesson: sometimes it is needed to look back, to look forward.

Competing interests

The authors declare that they have no competing interests.

References

- Adriano, A., 1932. Carmi Tradizioni Pregiudizi Nella Medicina Popolare Calabrese. Tipografia Pranno, Cosenza.
- Akkol, E.K., Güvenç, A., Yesilada, E., 2009. A comparative study on the antinociceptive and anti-inflammatory activities of five *Juniperus* taxa. *J. Ethnopharmacol.* 125, 330–336.
- Alberti, Á., Béni, S., Lackó, E., Riba, P., Al-Khrasani, M., Kéry, Á., 2012. Characterization of phenolic compounds and antinociceptive activity of *Sempervivum tectorum* L. leaf juice. *J. Pharm. Biomed. Anal.* 70, 143–150.
- Aliotta, G., Piomelli, D., Pollio, A., Touwaide, A., 2003. Le Pianta Medicinali del «Corpus Hippocraticum». Guerini e Associati, Milano.
- Atta, A.H., Abo EL-Sooud, K., 2004. The antinociceptive effect of some Egyptian medicinal plant extracts. *J. Ethnopharmacol.* 95, 235–238.
- Atta, A.H., Alkofahi, A., 1998. Anti-nociceptive and anti-inflammatory effects of some Jordanian medicinal plant extracts. *J. Ethnopharmacol.* 60, 117–124.
- Baeuerle, P.A., Henkel, T., 1994. Function and activation of NF- κ B in the immune system. *Annu. Rev. Immunol.* 12, 141–179.
- Beara, L.N., Orčić, D.Z., Lesjak, M.M., Mimica-Dukić, N.M., Peković, B.A., Popović, M.R., 2010. Liquid chromatography/tandem mass spectrometry study of anti-inflammatory activity of Plantain (*Plantago* L.) species. *J. Pharm. Biomed. Anal.* 52, 701–706.
- Begum, S., Saxena, B., Goyal, M., Ranjan, R., Joshi, V.B., Rao, C.V., Krishnamurthy, S., Sahai, M., 2010. Study of antiinflammatory, analgesic and antipyretic activities of seeds of *Hyoscyamus niger* and isolation of a new coumarinolignan. *Fitoterapia* 81, 178–184.
- Benincá, J.P., Dal Marco, J.B., Pizzolatti, M.G., Fröde, T.S., 2011. Analysis of the anti-inflammatory properties of *Rosmarinus officinalis* L. in mice. *Food Chem.* 124, 468–475.
- Bernoni, D.G., 1878. Tradizioni Popolari Veneziane. Filippi Editore, Venezia.
- Bertagnon, E., 1955. Sulla flora medicinale della Liguria. Usi tradizionali dell'Alta Fontanabuona. *Atti dell'Accademia Ligur. di Sci. e Lett.* 11, 201–214.
- Bitler, C.M., Viale, T.M., Damaj, B., Crea, R., 2005. Hydrolyzed olive vegetation water in mice Has anti-inflammatory activity. *J. Nutr.* 135, 1475–1479.
- Buenza, E.J., Schnepf, D.J., Bauer, B.A., Elkin, P.L., Riddle, J.M., Motley, T.J., 2004. Techniques: bioprospecting historical herbal texts by hunting for new leads in old tomes. *Trends Pharmacol. Sci.* 9, 494–498.
- Calixto, J.B., Otuki, M.F., Santos, A.R.S., 2003. Anti-inflammatory compounds of plant origin. Part I. action on arachidonic acid pathway, nitric oxide and nuclear factor κ B (NF- κ B). *Planta Med.* 69, 973–983.
- Calixto, J.B., Campos, M.M., Otuki, M.F., Santos, A.R.S., 2004. Anti-inflammatory compounds of plant origin. Part II Modulation of pro-inflammatory cytokines, chemokines and adhesion molecules. *Planta Med.* 70, 93–103.
- Calvia, G., 1927. Animali e piante nella tradizione popolare sarda e specialmente del Logudoro. *Il Folklore Italiano. Arch. Trimest. per la Racc. e lo Stud. delle Tradiz. Pop. Ital.* 11, 187–206.
- Calvo, M.I., 2006. Anti-inflammatory and analgesic activity of the topical preparation of *Verbena officinalis* L. *J. Ethnopharmacol.* 107, 380–382.
- Campêlo, L.M., de Almeida, A.A., de Freitas, R.L., Cerqueira, G.S., de Sousa, G.F., Saldanha, G.B., Feitosa, C.M., de Freitas, R.M., 2011. Antioxidant and antinociceptive effects of Citrus limon essential oil in mice. *J. Biomed. Biotechnol.* 1–8.
- Campillo, D., 1984. Neurosurgical pathology in prehistory. *Acta Neurochir.* 70, 275–290.
- Capuano, A., De Corato, A., Lisi, L., Tringali, G., Navarra, P., Dello Russo, C., 2009. Proinflammatory-activated trigeminal satellite cells promote neuronal sensitization: relevance for migraine pathology. *Mol. Pain.* 5, 43.
- Chan, K., Islam, M.W., Kamil, M., Radhakrishnan, R., Zakaria, M.N., Habibullah, M., Attas, A., 2000. The analgesic and anti-inflammatory effects of *Portulaca oleracea* L.

- subsp. *sativa* (Haw.) Celak. J. Ethnopharmacol. 73, 445–451.
- Chimenti, F., Cottiglia, F., Bonsignore, L., Casu, L., Casu, M., Floris, C., Secci, D., Bolasco, A., Chimenti, P., Granese, A., Befani, O., Turini, P., Alcaro, S., Ortuso, F., Trombetta, G., Loizzo, A., Guarino, I., 2006. Quercetin as the active principle of *Hypericum hircinum* exerts a selective inhibitory activity against MAO-A: extraction, biological analysis, and computational study. J. Nat. Prod. 69, 945–949.
- Chiovenda-Bensi, C., 1955. Piante medicinali nell'uso tradizionale della Valle d'Ossola. Atti dell'Accademia Ligur. di Sci. e Lett. 11, 32–52.
- Choi, E., Koo, S., 2005. Anti-nociceptive and anti-inflammatory effects of the ethanolic extract of potato (*Solanum tuberosum*). Food Agric. Immunol. 16, 29–39.
- Choi, E.M., Hwang, J.K., 2004. Antiinflammatory, analgesic and antioxidant activities of the fruit of *Foeniculum vulgare*. Fitoterapia 75, 557–565.
- Chou, S.C., Chiu, Y.J., Chen, C.J., Lin, Y.C., Wu, C.H., Chao, C.T., Chang, C.W., Peng, W.H., 2012. Analgesic and anti-inflammatory activities of the ethanolic extract of *Artemisia morrissonensis* Hayata in mice. Evid.-Based Complement. Altern. Med. 2012, 1–11.
- Choudhary, M.I., Jilil, S., Todorova, M., Trendafilova, A., Mikhova, B., Duddeck, H., Rahman, A., 2007. Inhibitory effect of lactone fractions and individual components from three species of the *Achillea millefolium* complex of Bulgarian origin on the human neutrophils respiratory burst activity. Nat. Product. Res. 21, 1032–1036.
- Chu, Y.F., Wise, M.L., Gulvady, A.A., Chang, T., Kendra, D.F., Jan-Willem, van Klinken, B., Shi, Y., O'Shea, M., 2013. In vitro antioxidant capacity and anti-inflammatory activity of seven common oats. Food Chem. 139, 426–431.
- Cirese, A.M., 1996. Lo studio delle tradizioni popolari. In: Brioschi, F., Di Girolamo, C. (Eds.), Manuale di Letteratura Italiana. Storia per Generi e Problemi. Dall'Unità d'Italia Alla Fine del Novecento Vol. IV. Bollati Boringhieri, Torino, pp. 921–941.
- Confessor, M.V.A., Mendonça, L.E.T., Mourão, J.S., Alves, R.R.N., 2009. Animals to heal animals: ethnoveterinary practices in semiarid region, Northeastern Brazil. J. Ethnobiol. Ethnomed. 5, 37.
- Conforti, F., Tundis, R., Marrelli, M., Menichini, F., Statti, G.A., De Cindio, B., Menichini, F., Houghton, P.J., 2010. Protective effect of *Pimpinella anisoides* ethanolic extract and its constituents on oxidative damage and its inhibition of nitric oxide in lipopolysaccharide-stimulated RAW 264.7 macrophages. J. Med. Food 13, 137–141.
- Coroneddi-Berti, C., 1877. Appunti di medicina Bolognese. In: Vol. I. (Ed.), Rivista di Letteratura Popolare. Ermanno Loescher Editore, Torino-Roma-Firenze, pp. 1–12.
- De Giacomo, G., 1896. Il Popolo di Calabria. Patitucci Editore, Castrovillari.
- De Nino, A., 1891. Usi e Costumi Abruzzesi. Tipografia G. Barbera, Firenze.
- De Vos, P., 2010. European materia medica in historical texts: longevity of a tradition and implications for future use. J. Ethnopharmacol. 132, 28–47.
- Díaz-Viciedo, R., Hortelano, S., Girón, N., Massó, J.M., Rodríguez, B., Villar, A., de Las Heras, B., 2008. Modulation of inflammatory responses by diterpene acids from *Helianthus annuus* L. Biochem. Biophys. Res. Commun. 369, 761–766.
- Domingues, R.B., Duarte, H., Rocha, N.P., Teixeira, A.L., 2015. Increased serum levels of interleukin-8 in patients with tension-type headache. Cephalalgia 35, 801–806.
- Drummond, E.M., Harbourn, N., Marete, E., Martyn, D., Jacquier, J., O'Riordan, D., Gibney, E.R., 2012. Inhibition of proinflammatory biomarkers in THP1 macrophages by polyphenols derived from chamomile, meadowsweet and willow bark. Phytother. Res. 27, 588–594.
- Erdemoglu, N., Küpeli, E., Yeşilada, E., 2003. Anti-inflammatory and antinociceptive activity assessment of plants used as remedy in Turkish folk medicine. J. Ethnopharmacol. 89, 123–129.
- Ferraro, G., 1885. Botanica popolare in Carpeneto d'Acqui. Arch. per lo Stud. delle Tradiz. Pop. IV, 165–189.
- Finamore, G., 1894. Tradizioni Popolari Abruzzesi. Carlo Clausen Editore, Torino-Palermo.
- Geraci, P.O., 1957. La medicina del popolo in Calabria. Folk. della Calabr. Riv. di Tradiz. Pop. I, 149–166.
- Gorji, A., 2003. Pharmacological treatment of headache using traditional Persian medicine. Trends Pharmacol. Sci. 7, 331–334.
- Green, M.W., Green, L.M., Rothrock, J.F., 2005. Headache for history buffs. In: Managing Your Headaches. Springer, New York, pp. 1–11.
- Guimarães, A.G., Xavier, M.A., de Santana, M.T., Camargo, E.A., Santos, C.A., Brito, F.A., Barreto, E.O., Cavalcanti, S.C., Antoniolli, A.R., Oliveira, R.C., Quintans-Júnior, L.J., 2012. Carvacrol attenuates mechanical hypernociception and inflammatory response. N-S Arch. Pharmacol. 385, 253–263.
- Háznagy-Radnai, E., Balogh, Á., Czige, S., Máthé, I., Hohmann, J., Blazsó, G., 2012. Antiinflammatory activities of Hungarian Stachys species and their iridoids. Phytother. Res. 26, 505–509.
- Headache Classification Committee of the International Headache Society, 2013. The international classification of headache disorders, 3rd edition (beta version). Cephalalgia 33 (9), 629–808.
- Hernández-Ortega, M., Ortiz-Moreno, A., Hernández-Navarro, M.D., Chamorro-Cevallos, G., Dorantes-Alvarez, L., Necoechea-Mondragón, H., 2012. Antioxidant, anti-nociceptive, and anti-inflammatory effects of carotenoids extracted from dried pepper (*Capsicum annuum* L.). J. Biomed. Biotechnol. 1–10.
- Huang, T.T., Lai, H.C., Chen, Y.B., Chen, L.G., Wu, Y.H., Ko, Y.F., Lu, C.C., Chang, C.J., Wu, C.Y., Martel, J., Ojcius, D.M., Chong, K.Y., Young, J.D., 2014. cis-Resveratrol produces anti-inflammatory effects by inhibiting canonical and non-canonical inflammasomes in macrophages. Innate Immun. Lond. 20, 735–750.
- Ibrahim, S.R., Mohamed, G.A., Al-Musayeb, N.M., 2012. New constituents from the rhizomes of Egyptian *Iris germanica* L. Molecules 17, 2587–2598.
- ISTAT, 2011. L'Italia in 150 Anni. Sommario di statistiche storiche 1861–2010. Istituto Nazionale di Statistica, Roma.
- Jacobo-Herrera, N.J., Vartiainen, N., Bremner, P., Gibbons, S., Koistinaho, J., Heinrich, M., 2006. NF-κB Modulators from Valeriana officinalis. Phytother. Res. 20, pp. 917–919.
- Jiang, J., Kang, T.B., Shim, D.W., Oh, N.H., Kim, T.J., Lee, K.H., 2013. Indole-3-carbinol inhibits LPS-induced inflammatory response by blocking TRIF-dependent signaling pathway in macrophages. Food Chem. Toxicol. 57, 256–261.
- Kaiser, P., Youssouf, M.S., Tasduq, S.A., Singh, S., Sharma, S.C., Singh, G.D., Gupta, V.K., Gupta, B.D., Johri, R.K., 2009. Anti-allergic effects of herbal product from *Allium cepa* (Bulb). J. Med. Food 12, 374–382.
- Karaca, M., Ozbek, H., Akkan, H.A., Tutuncu, M., Ozgokce, F., Him, A., Bakir, B., 2009. Anti-inflammatory activities of diethyl-ether extracts of *Helichrysum plicatum* DC. and *Tanacetum balsamita* L. in rats. Asian. J. Anim. Vet. Adv. 4, 320–325.
- Karenberg, A., Leitz, C., 2001. Headache in magical and medical papyri of ancient Egypt. Headache 21, 911–916.
- Karlsen, A., Paur, I., Böhn, S.K., Sakhi, A.K., Borge, G.I., Serafini, M., Erlund, I., Laake, P., Tonstad, S., Blomhoff, R., 2010. Bilberry juice modulates plasma concentration of NF-κB related inflammatory markers in subjects at increased risk of CVD. Eur. J. Nutr. 49, 345–355.
- Kim, B.H., Hong, S.S., Kwon, S.W., Lee, H.Y., Sung, H., Lee, I.J., Hwang, B.Y., Song, S., Lee, C.K., Chung, D., Ahn, B., Nam, S.Y., Han, S.B., Kim, Y., 2008. Diarctigenin, a lignan constituent from *Arctium lappa*, down-regulated zymosan-induced transcription of inflammatory genes through suppression of dna binding ability of nuclear factor-kappa B in macrophages. J. Pharmacol. Exp. Ther. 327, 393–401.
- Kosar, M., Küpeli, E., Malyer, H., Uylaser, V., Türkben, C., Baser, K.H., 2007. Effect of brining on biological activity of leaves of *Vitis vinifera* L. (Cv. Sultani C, ekirdeksiz) from Turkey. J. Agr. Food Chem. 55, 4596–4603.
- Kühn, C.G., 1826. Claudii Galeni Opera Omnia Tomus XII. Officina Libraria Car. Knoblochii, Lipsiae.
- Kühn, C.G., 1827. Claudii Galeni Opera Omnia Tomus XIV. Officina Libraria Car. Knoblochii, Lipsiae.
- Kühn, C.G., 1829. Pedanii Dioscoridis Anazarbei De Materia Medica Tomus I. Officina Libraria Car. Knoblochii, Lipsiae.
- Kumar, D., Kumar, S., Singh, J., Narender, Rashmi, Vashistha, B.D., Singh, N., 2010. Free radical scavenging and analgesic activities of *Cucumis sativus* L. Fruit. Extr. J. Young. Pharm. 2, 365–368.
- Laddomada, B., Durante, M., Minervini, F., Garbetta, A., Cardinali, A., D'Antuono, I., Caretto, S., Bianco, A., Mita, G., 2015. Phytochemical composition and anti-inflammatory activity of extracts from the whole-meal flour of Italian durum wheat cultivars. Int. J. Mol. Sci. 16, 3512–3527.
- Lanza, B., 2006. Cassano. Rimedi Popolari. In: il Regno delle due Sicilie descritto ed illustrato (Calabria. Ristampa anastatica). Edizioni Paparo, Napoli, pp. 1853–1860.
- Latronico, L., 1935. La Medicina infantile nelle superstizioni e nelle usanze popolaristiche italiane. Immacolata Concezione, Modena.
- Lee, H.G., Kim, H., Oh, W.K., Yu, K.A., Choe, Y.K., Ahn, J.S., Kim, D.S., Kim, S.H., Dinarello, C.A., Kim, K., Yoon, D.Y., 2004. Tetramethoxy hydroxyflavone p7F downregulates inflammatory mediators via the inhibition of nuclear factor kappa B. Ann. New Y. Acad. Sci. 1030, 555–568.
- Lee, I.S., Lim, J., Gal, J., Kang, J.C., Kim, H.J., Kang, B.Y., Choi, H.J., 2011. Anti-inflammatory activity of xanthohumol involves heme oxygenase-1 induction via NRF2-ARE signalling in microglial BV2 cells. Neurochem. Int. 58, 153–160.
- Lee, Y.C., Kim, S.H., Roh, S.S., Choi, H.Y., Seo, Y.B., 2007. Suppressive effects of *Chelidonium majus* methanol extract in knee joint, regional lymph nodes, and spleen on collagen-induced arthritis in mice. J. Ethnopharmacol. 112, 40–48.
- Leonti, M., Casu, L., Sanna, F., Bonsignore, L., 2009. A comparison of medicinal plant use in Sardinia and Sicily - De Materia Medica revisited? J. Ethnopharmacol. 121, 255–267.
- Lombardi-Satriani, R., 1970. Credenze popolari calabresi. Peloritana Editrice, Messina.
- Macêdo, S.B., Ferreira, L.R., Perazzo, F.F., Carvalho, J.C., 2004. Anti-inflammatory activity of *Arnica montana* 6cH: preclinical study in animals. Homeopathy 94, 84–87.
- Magiorkinis, E., Diamantis, A., Mitsikostas, D.D., Androustos, G., 2009. Headaches in antiquity and during the early scientific era. J. Neurol. 256, 1215–1220.
- Maham, M., Moslemzadeh, H., Jalilzadeh-Amin, G., 2013. Antinociceptive effect of the essential oil of tarragon (*Artemisia dracunculoides*). Pharm. Biol. 52, 208–212.
- Mancarella, A., 1930. Superstizioni popolari pugliesi. Il Folk. Ital. V, 126–151.
- Marzano, G.B., 1890. Usi e costumi di Laureana di Borrello - medicina popolare. La Calabr. III (4), 29–30.
- Mothana, R.A.A., 2011. Anti-inflammatory, antinociceptive and antioxidant activities of the endemic Soqotra *Boswellia elongata* Balf. f. and *Jatropha uniconostata* Balf. f. in different experimental models. Food Chem. Toxicol. 49, 2594–2599.
- Moussaid, M., Elamrani, A.E., Bourhim, N., Benaissa, M., 2011. In vivo anti-inflammatory and in vitro antioxidant activities of Moroccan medicinal plants. Nat. Product. Commun. 6, 1441–1444.
- Muhammad, I., Satoshi Takamatsu, S., Mustafa, J., Khan, S.I., Khan, I.A., Samoylenko, V., Mossa, J.S., El-Ferali, F.S., Dunbar, D.C., 2008. COX-2 inhibitory activity of Cafestol and analogs from coffee beans. Nat. Product. Commun. 3, 11–16.
- Nardi, G.M., 1935. La medicina popolare in Toscana. Lares VI, 272–282.
- Nasri, S., Anoush, M., Khatami, N., 2012. Evaluation of analgesic and anti-inflammatory effects of fresh onion juice in experimental animals. Afr. J. Pharm. Pharmacol. 6, 1679–1684.
- Orhan, D.D., Küpeli, E., Yeşilada, E., Ergun, F., 2006. Anti-inflammatory and antinociceptive activity of flavonoids isolated from *Viscum album* ssp. *Album*. Z. Nat. C., J. Biosci. 61, 26–30.
- Orlando, R.A., Gonzales, A.M., Hunsaker, L.A., Franco, C.R., Royer, R.E., Vander Jagt, D.L., Vander Jagt, D.J., 2010. Inhibition of nuclear factor κB activation and cyclooxygenase-2 expression by aqueous extracts of hispanic medicinal herbs. J. Med. Food 13, 1–4.
- Ostermann, V., 1894. La vita in Friuli. Usi, costumi, credenze, pregiudizi e superstizioni popolari Tipografia del Bianco Editore, Udine.
- Oztürk, N., Başer, K.H., Aydin, S., Oztürk, Y., Calış, I., 2002. Effects of *Gentiana lutea* spp

- symphyandra* on the central nervous system in mice. *Phytother. Res.* 16, 627–631.
- Pagano, L., 1992. *Natura Economia Storia in Calabria* Ristampa anastatica. Edizioni Cronaca di Calabria, Cosenza.
- Panico, A.M., Cardile, V., Garufi, F., Puglia, C., Bonina, F., Ronsisvalle, S., 2007. Effect of hyaluronic acid and polysaccharides from *Opuntia ficus indica* (L.) cladodes on the metabolism of human chondrocyte cultures. *J. Ethnopharmacol.* 111, 315–321.
- Park, E.H., Kahng, J.H., Lee, S.H., Shin, K.H., 2001. An anti-inflammatory principle from cactus. *Fitoterapia* 72, 288–290.
- Park, S.Y., Hong, S.S., Han, X.H., Hwang, J.S., Lee, D., Ro, J.S., Hwang, B.Y., 2007. Lignans from *Arctium lappa* and their inhibition of LPS-induced nitric oxide production. *Chem. Pharm. Bull.* 55, 150–152.
- Pasquarelli, G., 1987. *Medicina magia e classi sociali nella Basilicata degli anni venti. Scritti di un medico antropologo*. Cured by Bronzini, G.B., Congedo Editore, Galatina.
- Pedrotti, G., Bertoldi, V., 1930. *Nomi dialettali delle piante indigene del Trentino e della Ladinia Dolomitica*. Tipografia della Casa Editrice GB Monauini, Trento.
- Pépin, R., 1950. *Quintus Serenus Sammonicus, Liber Medicinalis*. Presses Universitaires de France, Paris.
- Pignatari, F., 1894. *Medicina popolare. Cefalaga-Coma*. La Calabr. 10, 76–77.
- Pitrè, G., 1896. *Medicina popolare siciliana*. Carlo Clausen Editore, Torino-Palermo.
- Plinio, G.S., 1985. *Storia Naturale*, vol. III*. Einaudi Editore, Torino.
- Pollio, A., De Natale, A., Appetiti, E., Aliotta, G., Touwaide, A., 2008. Continuity and change in the Mediterranean medical tradition: *ruta* spp. (rutaceae) in Hippocratic medicine and present practices. *J. Ethnopharmacol.* 116, 469–482.
- Potter, A., 1988a. *Hippocrates, Affections. Diseases I. Diseases II*. Harvard University Press, Cambridge (Massachusetts) and London.
- Potter, A., 1988b. *Hippocrates, Diseases III. Internal Affections. Regimen in Acute Diseases*. Harvard University Press, Cambridge (Massachusetts) and London.
- Raghav, S.K., Gupta, B., Shrivastava, A., Das, H.R., 2007. Inhibition of lipopolysaccharide-inducible nitric oxide synthase and IL-1 β through suppression of NF- κ B activation by 3-(1'-1'-dimethyl-allyl)-6-hydroxy-7-methoxy-coumarin isolated from *Ruta graveolens* L. *Eur. J. Pharmacol.* 560, 69–80.
- Ramadan, M., Goeters, S., Watzel, B., Krause, E., Lohmann, K., Bauer, R., Hempel, B., Imming, P., 2006. Chamazulene carboxylic acid and matricin: a natural proen and its natural produg, identified through similarity to synthetic drug substances. *J. Nat. Prod.* 69, 1041–1045.
- Rao, C.V., Verma, A.R., Gupta, P.K., Vijayakumar, M., 2007. Anti-inflammatory and antinociceptive activities of *Fumaria indica* whole plant extract in experimental animals. *Acta Pharm.* 57, 491–498.
- Ratheesh, M., Shyni, G.L., Sindhu, G., Helen, A., 2010. Protective effects of isolated polyphenolic and alkaloid fractions of *Ruta graveolens* L. on acute and chronic models of inflammation. *Inflammation* 33, 18–24.
- Redillas, C., Solomon, S., 2000. Prophylactic pharmacological treatment of chronic daily headache. *Headache* 40, 83–102.
- Reuter, U., Chiarugi, A., Bolay, H., Moskowitz, M.A., 2002. Nuclear factor-kappaB as a molecular target for migraine therapy. *Ann. Neurol.* 51, 507–516.
- Riddle, J.M., 1987. Folk tradition and folk medicine: recognition of drugs in classical antiquity. In: Scarborough, J. (Ed.), *Folklore and Folk Medicines*. American Institute of the History of Pharmacy, Madison, pp. 33–61.
- Rodrigues, M.R., Kanazawa, L.K., das Neves, T.L., da Silva, C.F., Horst, H., Pizzolatti, M.G., Santos, A.R., Baggio, C.H., Werner, M.F., 2012. Antinociceptive and anti-inflammatory potential of extract and isolated compounds from the leaves of *Salvia officinalis* in mice. *J. Ethnopharmacol.* 139, 519–526.
- Romeo, N., Gallo, O., Tagarelli, G., 2015. From disease to holiness: religious-based health remedies of Italian folk medicine (XIX-XX century). *J. Ethnobiol. Ethnomed.* 11, 50.
- Sapelli, G., 1991. *Dalla periferia all'integrazione europea, Storia dell'economia italiana. L'età contemporanea un paese nuovo*. Giulio Einaudi Editore, Torino.
- Sasannejad, P., Saeedi, M., Shoeibi, A., Gorji, A., Abbasi, M., Foroughipour, M., 2012. Lavender essential oil in the treatment of migraine headache: a placebo-controlled clinical trial. *Eur. Neurol.* 67, 288–291.
- Sayyah, M., Saroukhani, G., Peirovi, A., Kamalinejad, M., 2003. Analgesic and anti-inflammatory activity of the leaf essential oil of *Laurus nobilis* Linn. *Phytother. Res.* 17, 733–736.
- Sembianti, P., 1936. *Superstizione Nella Medicina Popolare Trentina*. Atti del III Congresso Nazionale di Arti e Tradizioni Popolari. Edizione dell'O.N.D., Roma.
- Shevel, E., 2011. The extracranial vascular theory of migraine-A great story confirmed by the facts. *Headache* 51, 409–417.
- SIISC, 2011. (Società Italiana per lo Studio delle Cefalee). *Linee guida per la terapia delle cefalee primarie*. Calzetti Mariucci Editore, Ferriera di Torgiano(PG).
- Speroni, E., Cervellati, R., Costa, S., Dall'Acqua, S., Guerra, M.C., Panizzolo, C., Utan, A., Innocenti, G., 2007. Analgesic and antiinflammatory activity of *Cyclamen repandum* S. et S. *Phytother. Res.* 21, 684–689.
- Stevens, P.F., 2001. *Angiosperm Phylogeny Website*. Version 12, July 2012. Available at <<http://www.mobot.org/MOBOT/research/APweb/>> Accessed 18th Feb 2016.
- Stratton, C.F., Newman, D.J., Derek, S., Tan, D.S., 2015. Cheminformatic comparison of approved drugs from natural product versus synthetic origins. *Bioorg. Med. Chem. Lett.* 25, 4802–4807.
- Strzelecka, M., Bzowska, M., Kozie, J., Szuba, B., Dubiel, O., Rivera Nunez, D., Heinrich, M., Berta, J., 2005. Anti-inflammatory effects of extracts from some traditional Mediterranean diet plants. *J. Physiol. Pharmacol.* 56, 139–156.
- Stulzer, H.K., Tagliari, M.P., Zampirolo, J.A., Cechinel-Filho, V., Schlemper, V., 2006. Antioedematogenic effect of marrubiin obtained from *Marrubium vulgare*. *J. Ethnopharmacol.* 108, 379–384.
- Sun, Y.X., Liu, J.C., Liu, D.Y., 2011. Phytochemicals and bioactivities of *anemone raddeana* Regel: a review. *Pharmazie* 66, 813–821.
- Tagarelli, G., Tagarelli, A., Piro, A., 2010. Folk medicine used to heal malaria in Calabria. *J. Ethnobiol. Ethnomed.* 6, 27.
- Tagarelli, G., Tagarelli, A., Liguori, M., Piro, A., 2013. Treating epilepsy in Italy between XIX and XX century. *J. Ethnopharmacol.* 145, 608–613.
- Tancredi, G., 1938. *Folclore garganico. Armillotta & Marino Tipografi, Manfredonia*.
- Thole, J.M., Kraft, T.F., Sueiro, L.A., Kang, Y.H., Gills, J.J., Cuendet, M., Pezzuto, J.M., Seigler, D.S., Lila, M.A., 2006. A comparative evaluation of the anticancer properties of European and American elderberry fruits. *J. Med. Food* 9, 498–504.
- Wang, Q., Kuang, H., Su, Y., Sun, Y., Feng, J., Guo, R., Chan, K., 2013. Naturally derived anti-inflammatory compounds from Chinese medicinal plants. *J. Ethnopharmacol.* 146, 9–39.
- WHO, 2011. *Atlas of Headache Disorders and Resources in the World*. WHO Press, Geneva, pp. 2011.
- Wittmann, D.H., Schein, M., Condon, R., 1996. Management of secondary peritonsillitis. *Ann. Surg.* 224, 10–18.
- Yesilada, E., Tanaka, S., Sezik, E., Tabata, M., 1988. Isolation of an anti-inflammatory principle from the fruit juice of *Ecballium elaterium*. *J. Nat. Prod.* 51, 504–508.
- Zakaria, Z.A., Gopalan, H.K., Zainal, H., Pojan, N.H.M., Morsid, N.A., Aris, A., Sulaiman, M.R., 2006. Antinociceptive, anti-inflammatory and antipyretic effects of *Solanum nigrum* chloroform extract in animal models. *Yakugaku Zasshi-J. Pharm. Soc. Jpn.* 126, 1171–1178.
- Zanchin, G., 2010. Headache: an historical outline. In: Finger, S., Boller, F., Tyler, K.L. (Eds.), *Handbook of Clinical Neurology. History of Neurology* 95. Elsevier, Amsterdam, pp. 375–386.
- Zanetti, Z., 1892. *La medicina delle nostre donne*. Lapiti Tipografo Editore, Città di Castello.